



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/435, A61K 38/17	A1	(11) International Publication Number: WO 97/33910 (43) International Publication Date: 18 September 1997 (18.09.97)
(21) International Application Number: PCT/US97/03753 (22) International Filing Date: 11 March 1997 (11.03.97) (30) Priority Data: 08/614,935 11 March 1996 (11.03.96) US (71) Applicant: THE ROCKEFELLER UNIVERSITY [US/US]; 1230 York Avenue, New York, NY 10021-6399 (US). (72) Inventor: KING, Te, Piao; 340 East 64th Street, New York, NY 10021 (US). (74) Agents: FEHLNER, Paul, F. et al.; Klauber & Jackson, 411 Hackensack Avenue, Hackensack, NJ 07601 (US).		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CU, CZ, EE, GE, GH, HU, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, TR, TT, UA, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: IMMUNOMODULATORY PEPTIDES OF VESPID ANTIGEN 5 (57) Abstract The present invention is directed to immunogenic peptides from vespid antigen 5. These immunogenic peptides can be used in immunotherapy for vespid venom allergic individuals. The present invention is thus directed to T cell epitopes of vespid antigen 5 that can anergize T cell responses in sensitive individuals.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

IMMUNOMODULATORY PEPTIDES OF VESPID ANTIGEN 5

The research leading to the present invention was supported by United States Public Health Service Grant No. AI-17021. The government may have certain rights in the invention.

5

FIELD OF THE INVENTION

The present invention is directed to immunogenic peptides from vespid antigen 5. These immunogenic peptides can be used in immunotherapy for vespid venom allergic individuals. The present invention is thus directed to T cell epitopes of vespid antigen 5 that can anergize T cell responses in sensitive individuals.

BACKGROUND OF THE INVENTION

15

Biochemical Aspects of Insect Venom Allergens

Insect sting allergy to bees and vespids is of common occurrence. The vespids include hornets, yellowjackets and wasps [Golden, et al., *Am. Med. Assoc.*, **262**:240 (1989)]. Susceptible people can be sensitized on exposure to minute amounts of venom proteins; as little as 2-10 μ g of protein is injected into the skin on a single sting by a vespid [Hoffman and Jacobson, *Ann. Allergy.*, **52**:276 (1984)].

Indeed, venom allergens from insects of the Hymenoptera order have been extensively studied. These insects are bees, vespids and fire ants. The bees include honey bees (*Apis mellifera*) and bumble bees (*Bombus pennsylvanicus*). The vespids include hornets (*Dolichovespula spp.*; *Vespa crabo*), yellow jackets (*Vespula spp.*), and paper wasps (*Polistes spp.*). In Table 1 are listed the venom allergens from these insects with known primary structures.

25

Table 1

Cloned and/or sequenced insect venom allergens

	Allergen <u>name¹</u>	<u>Common name</u>	Mol. <u>Size²</u>	<u>Recombinant protein³</u> <u>unfolded</u> <u>folded</u>	
5					
		Bumble bee, <i>Bombus pennsylvanicus</i>			
	Bom p 1	phospholipase A ₂	16 kd		
	Bom p 4	protease	28 kd	—	—
10					
		Honey bee, <i>Apis mellifera</i>			
	Api m 1	phospholipase A ₂	16 kd	+	+
	Api m 2	hyaluronidase	39 kd	+	+
	Api m 3	acid phosphatase	43 kd		
	Api m 4	melittin	3 kd	—	—
15					
		Fire ant, <i>Solenopsis invicta</i>			
	Sol i 1	phospholipase A ₁	37 kd	—	—
	Sol i 2		30 kd	—	—
	Sol i 3	antigen 5	23 kd	—	+
20	Sol i 4		20 kd	—	—
		White face hornet, <i>Dolichovespula maculata</i>			
	Dol m 1	phospholipase A ₁	34 kd	+	—
	Dol m 2	hyaluronidase	38 kd	+	—
25	Dol m 5	antigen 5	23 kd	+	+
		European hornet, <i>Vespa crabro</i>			
	Vesp c 1	phospholipase A ₁	23 kd	—	—
	Vesp c 5	antigen 5	23 kd	—	—
30					
		Paper wasp, <i>Polistes annularis</i>			
	Pol 1 5 ⁵	antigen 5	23 kd	+	—
		Yellow jacket, <i>Vepula vulgaris</i>			
35	Ves v 1	phospholipase A ₁	34 kd	+	—
	Ves v 2	hyaluronidase	38 kd	+	—
	Ves v 5 ⁵	antigen 5	23 kd	+	—

Footnotes

- Allergen names are designated according to an accepted nomenclature system [King et al., *WHO Bulletin*, **72**:797 (1994)].
- Several allergens are glycoproteins, and the molecular size given refers only to the protein portion.
- + and - signs refer to the availability of recombinant proteins.
- Sequence of antigen 5 from *S. richteria* is known [Smith & Hoffman, *J. Allerg. Clin. Immunol.*, **89**:293 (1992)].
- Sequences of antigen 5s from several other vespids are known; *D. arenaria* [Lu et al., *J. Immunol.* **150**:2823 (1993)], *P. exclamans* and *P. fuscatus*, and *V. flavopilosa*, *V. germanica*, *V. maculifrons*, *V. pennsylvanica*, *V. spumosa* and *V. vidua* [Hoffman et al., *Int. Archs. Allergy App. Immunol.*, **84**:24 (1987)].

- There are many species of hornets (genus *Dolichovespula*), yellowjackets (genus *Vespula*) and wasp (genus *Polistes*) in North America [Akre, et al., "Yellowjackets of America North of Mexico," Agriculture Handbook No. 552, US Department of Agriculture (1980)]. The vespids have similar venom compositions [King, et al., *Biochemistry*, 5 17:5165 (1978); King, et al., *Mol. Immunol.* 20:297 (1983); King, et al., *Arch. Biochem. Biophys.* 230:1 (1984); King, et al., *J. Allergy and Clin. Immunol.*, 75:621 (1985); King, *J. Allergy Clin. Immunol.*, 79:113 (1987); Hoffman, *J. Allergy and Clin. Immunol.*, 75:611 (1985)]. Their venom each contains three major venom allergens, phospholipase (37 kD), hyaluronidase (43 kD) and antigen 5 (23 kD) of as yet unknown biologic 10 function. Homologous venom allergens from hornets, yellow jackets, and paper wasps have high degrees of sequence identity ranging from about 70% for antigen 5s to about 90% for hyaluronidases [Lu et al., *J. Immunol.*, 150:2823 (1993)].
- 15 Antigen 5s from several species each of hornets, yellowjackets and paper wasps have been cloned and/or sequenced [Fang et al., *Proc. Natl. Acad. Sci., USA*, 85:895-899 (1988); Lu et al., *supra*; Hoffman, *J. Allergy Clin. Immunol.*, 92:707-716 (1993)]. For phospholipases and hyaluronidases only those from hornets and yellowjackets have been cloned and/or sequenced [Soladatova et al., *FEBS Letters*, 320:145-149 (1993); Lu et al., 20 *J. Biol. Chem.*, 270:4457-4465 (1995); King et al., *J. Allergy Clin. Immunol.*, (1996); Hoffman, *Int. Arch. Allergy Immunol.*, 104:184-190 (1994)]. One common feature of these venom proteins is their varying extent of sequence homology with mammalian proteins.
- 25 White faced hornet (*Dolichovespula maculata*) has three forms of antigen 5. Two of these forms, Dol m 5.01 and 5.02, differ in 23% of their sequences, and they are antigenically cross reactive at both B and T cell levels [Fang et al., *Proc. Natl. Acad. Sci., USA*, 85:895-899 (1988); Lu et al., *J. Immunol.*, 150:2823-2830 (1993)]. The studies described here were made with Dol m 5.02, also referred to as hornet Ag5 form 2. The amino acid 30 sequences of Dol m 5.01 and 5.02, as well as those of the homologous antigen 5s from yellow hornet (*Dolichovespula arenaria*), two species of yellowjacket (*Vespula maculifrons* and *vulgaris*) and two species of papers wasps (*Polistes annularis* and *exclamans*) are given in Figure 1.

Fire ant venom contains four allergens. They are antigen 5, phospholipase A₁, and Sol i 2 and 4. Fire ant antigen 5 has about 50% sequence identity with vespid antigen 5 [Hoffman, *J. Allergy Clin. Immunol.*, **91**:71 (1995)]. Only partial sequence data is available for fire ant phospholipase, and it shows sequence identity with vespid phospholipase A₁ [Hoffman, *J. Allergy Clin. Immunol.*, **95**:372 (1995)].

Bumble bee venom has two allergens of known sequences; phospholipase A₂ and protease. But honey bee venom has four allergens of known sequences; acid phosphatase, phospholipase A₂, hyaluronidase and a cytolytic peptide melittin. The two bee venom phospholipase A₂ have extensive sequence identity and they are not-related to vespid phospholipase A₁ [Hoffman, "Hymenoptera Venom Proteins" in *National Toxins*, R.B. Singh (ed.), Plenum Publishing 6 (1996)]. Honey bee venom hyaluronidase has about 55% sequence identity with the homologous vespid hyaluronidases.

In addition to the insect venom allergens described above, the complete amino acid sequence of several major allergens from different grass [Perez, et al., *J. Biol. Chem.*, **265**:16210 (1990); Ansari, et al., *Biochemistry*, **26**:8665 (1989); Silvanovich, et al., *J. Biol. Chem.*, **266**:1204 (1991)], tree pollen [Breiteneder, *EMBO J.*, **8**:1935(1989); Valenta, et al., *Science*, **253**:557 (1991)], weed pollen [Rafnar, et al., *J. Biol. Chem.*, **266**:1229 (1991); Griffith, et al., *Int. Arch. Allergy Appl. Immunol.*, **96**:296 (1991)], mites [Chua, et al., *J. Exp. Med.*, **167**:175 (1988)], cat dander [Griffith, et al., *Gene*, **113**:263 (1992)], and mold [Aruda, et al., *J. Exp. Med.*, **172**:1529 (1990); Han, et al., *J. Allergy Clin. Immunol.*, **87**:327 (1991)] have been reported in the past few years. These major allergens are proteins of 10-40 kD and they have widely different biological functions. Nearly all allergens of known sequences have a varying extent of sequence similarity with other proteins in our environment.

T and B Cell Epitopes of Allergens

Antibody responses to proteins require the collaboration of T helper and B lymphocytes and antigen presenting cells (APC). The antigen receptors of B cells are the membrane-bound antibody (Ab) molecules, which recognize and bind immunogens directly. The antigen receptors of T cells (TCR) only recognize and bind complexes of antigenic peptide-MHC class II molecule. Immunogens are first processed by APC into peptides that are presented on the surface of APC in association with the MHC class II molecules

[Unanue, *Current Opinion in Immunol*, 4:63 (1992)]. As MHC molecules are highly polymorphic in individuals, they have different specificity of binding antigenic peptides [Rothbard and Gefer, *Ann. Rev. Immunol.*, 9:527 (1991)]. This is one mechanism for genetic control of immune response.

5

T helper cells are activated when the antigen receptor binds the peptide-MHC complex on the surface of APC. Activated T cells secrete lymphokines. In mice [Street and Mosmann, *FASEB J.*, 5:171 (1991)] and apparently in humans [Wierenga, et al., *J. Immunol.*, 144:4651 (1990); Parronchi, et al., *Proc. Natl. Acad. Sci. USA.*, 88:4538 (1991)] the T helper cells can be divided into different types on the basis of their patterns of lymphokine production. Primarily, T helper cells divide into two groups: TH1 cells producing IL-2 and IFN- γ , and TH2 cells producing IL-4 and IL-5. These lymphokines in turn influence the antigen-activated B cells to differentiate and proliferate into plasma cells secreting Abs of different isotypes. IL-4 is one lymphokine known to influence IgE synthesis [Finkelman, et al., *Ann. Rev. Immunol.*, 8:303 (1990)].

It is believed that the entire accessible surface of a protein molecule can be recognized as epitopes by the antigen receptors of B cells, although all epitopes are not necessarily recognized with equal likelihood [Benjamin, et al., *Ann. Rev. Immunol.*, 2:67 (1984)]. B cell epitopes of a protein are of two types: topographic and linear. The topographic type consists of amino acid residues which are spatially adjacent but may or may not be sequentially adjacent. The linear type consists of only sequentially adjacent residues. X-ray crystallographic data of Ag-Ab complexes indicate the size of their complementary binding region to have 16-17 amino acid residues [Amit, et al., *Science*, 233:747 (1986)], but peptide mapping suggests that less than about 8 residues contribute significantly to the binding process of a linear epitope [Appel, et al., *J. Immunol.*, 144:976 (1990)].

Allergens, like other protein antigens, can have both types of B cell epitopes or only one. For example, vespid antigen 5s have both types [King et al., *J. Immunol.*, 154:577 (1995)]. Bee venom melittin appears to have only one B cell epitope of linear type [King, et al., *J. Immunol.*, 133:2668 (1984)].

T cell epitopes of proteins consist of only the linear type since they are peptides that have been processed in the lysosomes of APC by proteases of unknown specificity [Unanue,

Curr. Op. Immunol., **4**:63 (1992)]. Analysis of naturally processed antigenic peptides bound to MHC class II molecules indicates that their size ranges from about 13 to 17 amino acid residues, but analysis of synthetic peptide-MHC class II molecule complex for their T cell proliferate response suggests a minimal size of about 8 amino acid residues [Cf. Rudensky et al., *Nature*, **353**:622 (1991)]. Studies suggest that T cell epitopes are distributed throughout the entire protein molecule, and they may function as major or minor determinants depending on the MHC haplotype of the immunized host [Roy, et al., *Science*, **244**:572; Gammon, et al., *Immunol. Rev.*, **98**:53 (1987); O'Hehir et al., *Ann. Rev. Immunol.*, **9**:67 (1991)].

10

Hypersensitivity of the immediate type is known to be caused by the presence of allergen-specific IgE. IgE is found in the circulation and bound to specific IgE-Fc receptors on mast cells and basophils. Cross-linking of cell-bound IgE by allergens leads to release of histamine, leukotrienes and other chemical mediators that cause the allergic symptoms.

15 IgE is one of the different isotypes of immunoglobulins. As pointed out above, lymphokines secreted by T cells influence isotype switch events in B cells.

Because of the central role of TH2 cells in determining the isotypes switch event of B cells, the T cell epitopes of several allergens have been mapped [Cf. O'Hehir et al., *supra*]. These allergens include ragweed *Amb a* III, rye grass *Lol p* I, cat *Fel d* I, mouse urine *Mus m* I, midge *Chi t* I, bee venom phospholipase A₂ [Dhillon, et al., *J. Allergy Clin. Immunol.*, **90**:42 (1992)] melittin [Fehlner, et al., *J. Immunol.*, **146**:799 (1991)], and hornet antigen 5 [King et al., *J. Allergy Clin. Immunol.*, **91**:283 (1993)]. The data do not reveal any unusual or common structural features. However, any conclusion from these data is qualified as these data are collected from humans and mice of different haplotypes.

25

Modulation of T and B Cell Responses

Normally hosts are tolerant to the dominant B and T cell epitopes of self proteins by clonal deletion and anergy. However this tolerance can be broken under certain circumstances [Gammon, et al., *Immunol. Today.*, **12**:193 (1991); Basten, et al., *Immunol. Rev.*, **122**:5 (1991)]. It has been suggested that self-tolerance is broken in autoimmune diseases through encounters with foreign proteins that are similar to host proteins. Therefore the sequence similarity of allergens with autologous proteins is of interest for closer investigation.

30

Mature B cells are activated in response to multi-valent antigens which can cross-link cell surface Ig receptors [DeFranco, *Ann. Rev. Cell Biol.*, **3**:143 (1987)], and they are rendered anergic in response to mono-valent antigen [Basten, et al., 1991, *supra*]. Antigen activation of T cells requires not only the integration of TCR with peptide-MHC complex but also with other co-stimulating signals on the surface of APC [Schwartz, *Science*, **248**:1349 (1990); Jenkins and Miller, *FASEB J.*, **6**:2428 (1992)]. Interaction of TCR with peptide-MHC complex in absence of co-stimulating signals can lead to T cell anergy.

- 10 The molecular mechanism of B or T cell anergy is not yet understood [Cf. Schwartz, 1990, *supra*; Jenkins and Miller, 1992, *supra*; Ales-Martinez, et al., *Immunol. Today*, **12**:201 (1991)]. *In vitro* studies with T cell clones reveals that occupancy of TCR by artificial peptide-MHC complex in absence of co-stimulating signals leads to altered intracellular signal transduction and/or repressor gene activation which can prevent lymphokine transcription.

- Early studies have shown that the physical state of the immunogen and the route of immunization are important variables in determining the outcome of an immune response. In the light of our current understanding, these variables may well influence antigen presentation so as to have T and B cell activation or anergy.

- One way to treat allergic diseases is by immunotherapy which involves repeated subcutaneous injections of the offending allergen(s) into patients. The amounts of allergens which can be injected are limited by the danger of unwanted systemic allergic reaction in patients. For most patients following immunotherapy, their allergen-specific IgE levels initially rise followed with gradual decrease of their allergen-specific IgE levels, and there is also down-regulation of allergen-specific T cell responses [P.S. Norman, *Current Op. Immunol.*, **5**:968 (1993)].

- 30 Because of the undesirable systemic reaction on immunotherapy with native allergens, there has been continued interest in the development of modified allergens with reduced allergenic activities for immunotherapy [T.P. King, in "Bronchial Asthma," edited by E.B. Weiss and M. Stein, Little Brown, Boston, pp. 43-49 (1993); R.E. O'Hehir et al., 1991, *supra*].

Three reports have appeared recently on the use of T cell epitope peptides to modulate allergen-specific immune responses. One report is on the subcutaneous injection of mice with two peptides from the major cat allergen Fel d I to decrease T cell response to the entire molecule Fel d I [Briner et al., *Proc. Natl. Acad. Sci. U.S.A.*, **90**:7608-12 (1993)].

- 5 Another is on the intranasal therapy with a peptide from the major mite allergen Der p I to suppress allergen-specific response in naive or sensitized mice [Hoyne et al., *J. Exp. Med.*, **178**:1783-1788 (1993)]. The third reports that peptides containing T cell epitopes of bee venom phospholipase A₂ were used successfully in immunotherapy to treat patients with bee venom allergy [Müller et al., *J. Allergy Clin. Immunol.* **97**:426 (1996)].

10

Since an MHC class II molecule of any one haplotype can bind a wide range of peptides in its binding groove, it may be possible to modulate T cell response by inhibition of allergen-derived T cell epitope binding to MHC molecules with other peptides. For example, a mouse lysozyme peptide which is not immunogenic by itself in H-2^k mice

15 inhibits T cell response to hen egg white lysozyme [Adorini and Nagy, *Immunol. Today.*, **11**:21 (1990)]. Another example is the *in vitro* inhibition of T cell response to a mite allergen by an influenza HA peptide [O'Hehir et al., *J. Allergy Clin. Immunol.*, **87**:1120 (1991)].

- 20 Experimental autoimmune encephalomyelitis (EAE) in mice or rats is a well-studied model for multiple sclerosis. Many studies have identified immunodominant T cell determinants for myelin basic protein, which is used to induce this condition. Peptides that correspond to immunodominant epitopes of myelin basic protein can induce tolerance to the same peptide antigen or to the intact myelin basic protein. The same peptides that induced
- 25 tolerance could also induce T cell anergy in an ongoing autoimmune response [Gaur et al., *Science*, **259**:1491-1494 (1992)].

It has been reported that subcutaneous or intranasal pretreatment of mice with T cell

epitope peptides of the major cat allergen Fel d 1 [Briner et al., *Proc. Natl. Acad. Sci.*

- 30 *USA*, **90**:7608-7612 (1993)] or the major mite allergen Der p 1 [Hoyne et al., *J. Exp. Med.*, **178**:1783-1788 (1993)] lead to T cell anergy and that reduced antibody responses are observed on subsequent immunization with the allergen. T cell epitope peptides of Fel d 1 studied in the murine system are currently being evaluated for immunotherapy of patients [Norman et al., *J. Allergy Clin. Immunol.*, **95**:259 (1995)]. Published data

indicate that the T cell epitope regions of Der p 1 detected in the murine system overlap those found in humans [O'Hehir et al., *Europ. J. Clin. Invest.*, **23**:763 (1993)]. On the basis of findings with Fel d 1 and Der p 1, it is reasonable to conclude that the T cell epitope data of white faced hornet Ag5 obtained in mice as described herein will be applicable for studies in humans.

There remains a need in the art for the identification of T cell epitopes of vespid venom allergens, particularly antigen 5, for immunotherapy of venom allergy.

There is also a need in the art to use peptides having T cell epitopes of vespid venom allergens to study induction of tolerance in mice and induction of tolerance in humans.

There is a further need to test whether a modified peptide inhibits allergen T cell epitope binding to MHC class II molecule, or induces T cell anergy, or both.

These and other needs in the art are satisfied by the present invention.

The citation of references herein shall not be construed as an admission that such is prior art to the present invention.

SUMMARY OF THE INVENTION

The present invention provides the sequences of immunodominant peptides of vespid venom antigen 5, and corresponding peptides from other antigens, such as fire ant Sol i 3.

An immunodominant peptide is one that contains a T cell epitope of the antigen, such that T cells immunized with the antigen will be stimulated when contacted with the peptide. Such peptides of the invention are preferably immunomodulatory peptides as well, in that they induce T cell anergy when administered to a subject, or otherwise affect the immune response of the subject.

In yet another embodiment, the present invention provides a pharmaceutical composition effective for the treatment of a vespid venom allergen-specific allergic condition comprising a polypeptide of the invention that has an immunomodulatory portion of a T cell epitope of a vespid venom antigen 5. More particularly, the invention provides

pharmaceutical compositions comprising such polypeptides, including different isoforms, of a vespid venom antigen 5, for example, *Dolichovespula maculata*, *Vespula vulgaris*, *Vespula maculifrons*, *Dolichovespula arenaria*, *Polistes annularis*, and *Polistes exclamans*.

- 5 In yet still another embodiment, the present invention provides a method for treating a vespid venom allergen-specific condition comprising administering a therapeutically effective dose of a pharmaceutical composition of the invention.

In its broadest aspect, the present invention is directed to a peptide characterized by
 10 having between 8 (the generally recognized minimum number of amino acids for an antigenic T cell epitope) and 35 amino acid residues of vespid venom antigen 5; and being antigenic for T cell proliferation in a mouse immunized with a vespid venom antigen 5, which mouse is a strain selected from the group consisting of BALB/c, ASW/Sn, C3H/He, and P/J. More particularly, a peptide of the invention corresponds to a fragment
 15 of white face hornet antigen 5, form 2, selected from the group consisting of:

	NNYCKIKCRKGIHTLCKFGT	(SEQ ID NO:8);
	GIHTLCKFGTSMKPNCGRNV	(SEQ ID NO:9);
	KNEILKRHNDFRQNVAKGLE	(SEQ ID NO:12);
	FRQNVAKGLETRGKPGPQPP	(SEQ ID NO:13);
20	TRGKPGPQPPAKNMNVLVWN	(SEQ ID NO:14);
	DELAQIAQTWANQCDFNHDD	(SEQ ID NO:16);
	CRNTAKYQVGQNIAISSTTA	(SEQ ID NO:18);
	QNIAISSTTATQFDRPSKLI	(SEQ ID NO:19);
	TQFDRPSKLIKQWEDEVTEF	(SEQ ID NO:20);
25	KQWEDEVTEFNYKVGLQNSN	(SEQ ID NO:21);
	NYKVGLQNSNFRKVGHYTQM	(SEQ ID NO:22);
	FRKVGHYTQMVWGKT	(SEQ ID NO:23);
	KEIGCGSIKYIEDNWYTHYL	(SEQ ID NO:25); and
	IEDNWYTHYLVLCNYGPGGND	(SEQ ID NO:26).

- 30 Such peptides may be from the specific antigen, white face hornet antigen 5, form 2, or may be from the corresponding (homologous) segments of other vespid venom antigens, such as but not limited to *Dolichovespula maculata*, *Vespula vulgaris*, *Vespula maculifrons*, *Dolichovespula arenaria*, and *Polistes exclamans*. Other sources of antigen 5 of the invention include *Vespa crabro* (European hornet), *V. flavopilosa* (yellow jacket), *V.*

germanica (yellowjacket), *V. pennsylvanica* (yellowjacket), *V. squamosa* (yellowjacket), *V. vidua* (yellowjacket), and *P. fuscatus* (paperwasp), or fire ant allergen Sol i 3, which shares sequence similarity with (and possibility is homologous to) vespid venom antigen 5s. Such corresponding peptides from various species are referred to herein as

5 homologues, as they represent variants from the corresponding, *i.e.*, homologous, alleles of the antigen 5 gene from different vespid species. The term allelic variant includes, as well, single amino acid variations in antigen 5s from the same species, including different isoforms (such as the three isoforms found for white face hornet antigen 5), and mutations.

10

Preferably, the peptide corresponds to a fragment of white face hornet antigen 5, form 2, or a homologue thereof, selected from the group consisting of:

	NNYCKIKCRKGIHTLCKFGT	(SEQ ID NO:8);
	GIHTLCKFGTSMKPNCGRNV	(SEQ ID NO:9);
15	KNEILKRHNDFRQNVAKGLE	(SEQ ID NO:12);
	FRQNVAKGLETRGKPGPQPP	(SEQ ID NO:13);
	DELAQIAQWTWQNCDFNHDD	(SEQ ID NO:16);
	NYKVGLQNSNFRKVGHYTQM	(SEQ ID NO:22);
	FRKVGHYTQMVGWKT	(SEQ ID NO:23); and
20	IEDNWYTHYLVLCNYGPGGND	(SEQ ID NO:26).

This group of peptides showed significant T cell stimulation of antigen 5-immunized T cells in more than one species.

As pointed out above, the present invention advantageously provides not only the
 25 immunodominant T cell epitopes for white face hornet antigen 5, but corresponding epitopes from antigen 5s from other vespid species, particularly subfamilies Vespinae and Polistinae, more particularly the genera *Vespa*, *Vespula*, *Dolichovespula*, and *Polistes*. In addition, the invention provides for consensus polypeptides, which incorporate the same amino acid residues at conserved positions, and any amino acid residue or amino acid
 30 residues having similar properties at divergent positions. In other words, the present invention contemplates immunodominant peptides that incorporate polymorphisms found among the antigen 5 homologues from different species.

In a specific embodiment, the peptide has the amino acid sequence

X_1 BYCKIX₂CX₃X₄GX₅X₆HTX₇CX₈X₉G (SEQ ID NO:31), wherein X_1 is a neutral amino acid residue, X_2 is a basic amino acid residue or deleted, X_3 is any amino acid residue, X_4 is a polar amino acid, X_5 is glycine or deleted, X_6 is any amino acid residue, X_7 is an amino acid residue with an aliphatic side chain, X_8 is a polar amino acid residue, and X_9 is an amino acid residue with an aromatic side chain. In a more specific embodiment, X_1 is valine or asparagine; X_2 is lysine, arginine, or deleted; X_3 is proline, arginine, serine, or leucine; X_4 is arginine, lysine, or serine; X_6 is isoleucine, leucine, threonine, or valine; X_7 is alanine, valine, isoleucine, or leucine; X_8 is lysine, arginine, glutamine, or asparagine; or X_9 is phenylalanine, tryptophan, or tyrosine; or any combination of the foregoing.

In another specific embodiment, the peptide has the amino acid sequence

GX₅X₆HTX₇CX₈X₉GX₁₀SX₁₁KPX₁₂X₁₃NCX₁₄X₁₅X₁₆X₁₇X₁₈ (SEQ ID NO:32), wherein X_5 is glycine or deleted, X_6 is any amino acid residue, X_7 is an amino acid residue with an aliphatic side chain, X_8 is a polar amino acid residue, X_9 is an amino acid residue with an aromatic side chain, X_{10} is a polar amino acid residue, X_{11} is any amino acid residue, X_{12} is a polar amino acid residue or deleted, X_{13} is a basic amino acid residue or deleted, X_{14} is a small chain amino acid residue, X_{15} is any amino acid residue, X_{16} is a basic or polar neutral amino acid residue, X_{17} is any amino acid residue, and X_{18} is an aliphatic amino acid residue. More specifically, X_6 is isoleucine, leucine, threonine, or valine; X_7 is alanine, valine, isoleucine, or leucine; X_8 is lysine, arginine, glutamine, or asparagine; X_9 is phenylalanine, tryptophan, or tyrosine; X_{10} is threonine, serine, aspartic acid, or glutamic acid; X_{11} is leucine, isoleucine, methionine, or threonine; X_{12} is serine or deleted; X_{13} is lysine, arginine, or deleted; X_{14} is glycine or alanine; X_{15} is asparagine, glutamine, glycine, serine, arginine, or lysine; X_{16} is lysine, arginine, asparagine, or glutamine; X_{17} is lysine, arginine, isoleucine, leucine, or valine; or X_{18} is alanine, valine, isoleucine, or leucine; or any combination of the foregoing.

The present invention also advantageously provides for combining two overlapping peptides into a larger peptide having up to 35 amino acid residue. In one such embodiment, the peptide has the amino acid sequence

X_1 BYCKIX₂CX₃X₄GX₅X₆HTX₇CX₈X₉GX₁₀SX₁₁KPX₁₂X₁₃NCX₁₄X₁₅X₁₆X₁₇X₁₈ (SEQ ID NO:33), wherein X_1 is a neutral amino acid residue, X_2 is a basic amino acid residue or

deleted, X₃ is any amino acid residue, X₄ is a polar amino acid, X₅ is glycine or deleted, X₆ is any amino acid residue, X₇ is an amino acid residue with an aliphatic side chain, X₈ is a polar amino acid residue, X₉ is an amino acid residue with an aromatic side chain, X₁₀ is a polar amino acid residue, X₁₁ is any amino acid residue, X₁₂ is a polar amino acid residue or deleted, X₁₃ is a basic amino acid residue or deleted, X₁₄ is a small chain amino acid residue, X₁₅ is any amino acid residue, X₁₆ is a basic or polar neutral amino acid residue, X₁₇ is any amino acid residue, and X₁₈ is an aliphatic amino acid residue. In a more specific embodiment, X₁ is valine or asparagine; X₂ is lysine, arginine, or deleted; X₃ is proline, arginine, serine, or leucine; X₄ is arginine, lysine, or serine; X₅ is isoleucine, leucine, threonine, or valine; X₆ is alanine, valine, isoleucine, or leucine; X₇ is alanine, valine, isoleucine, or leucine; X₈ is lysine, arginine, glutamine, or asparagine; X₉ is phenylalanine, tryptophan, or tyrosine; X₁₀ is threonine, serine, aspartic acid, or glutamic acid; X₁₁ is leucine, isoleucine, methionine, or threonine; X₁₂ is serine or deleted; X₁₃ is lysine, arginine, or deleted; X₁₄ is glycine or alanine; X₁₅ is asparagine, glutamine, glycine, serine, arginine, or lysine; X₁₆ is lysine, arginine, asparagine, or glutamine; X₁₇ is lysine, arginine, isoleucine, leucine, or valine; or X₁₈ is alanine, valine, isoleucine, or leucine; or any combination of the foregoing.

In another specific embodiment, the peptide has the amino acid sequence KX₁₉X₂₀IX₂₁X₂₂X₂₃HNX₂₄FRQKX₂₅AX₂₆GLE (SEQ ID NO:34), wherein X₁₉ is a basic or neutral polar amino acid residue X₂₀ is any amino acid residue X₂₁ is an aliphatic amino acid residue, X₂₂ is a polar amino acid residue, X₂₃ is a polar charged amino acid residue, X₂₄ is a polar amino acid residue, X₂₅ is an aliphatic amino acid residue; and X₂₆ is a polar basic or neutral amino acid residue. More specifically, X₁₉ is asparagine, glutamine, or lysine; X₂₀ is aspartic acid, glutamic acids, leucine, or isoleucine; X₂₁ is valine, leucine, or isoleucine; X₂₂ arginine, asparagine, or serine; X₂₃ is glutamic acid or arginine; X₂₄ is aspartic acid, glutamic acid, asparagine, glutamine, or arginine; X₂₅ is valine, leucine, or isoleucine; or X₂₆ is arginine, asparagine, or lysine; or any combination of the foregoing.

In still another specific embodiment, the peptide has the amino acid sequence FRQKX₂₅AX₂₆GLETRGX₂₇PGPQPX₂₈ (SEQ ID NO:35), wherein X₂₅ is an aliphatic amino acid residue, X₂₆ is a polar basic or neutral amino acid residue, X₂₇ is a polar basic or neutral amino acid residue, and X₂₈ is any amino acid residue. More specifically, X₂₅ is valine, leucine, or isoleucine; X₂₆ is arginine, asparagine, or lysine; X₂₇ is asparagine or lysine; or X₂₈ is glycine, alanine, or proline; or any combination of the foregoing.

As noted above, overlapping peptides may be provided as a single, larger peptide. Thus, in another embodiment, the peptide has the amino acid sequence

$KX_{19}X_{20}IX_{21}X_{22}X_{23}HNX_{24}FRQKX_{25}AX_{26}GLETRGX_{27}PGPQPX_{28}$ (SEQ ID NO:36), wherein

X_{19} is a basic or neutral polar amino acid residue X_{20} is any amino acid residue X_{21} is an
 5 aliphatic amino acid residue, X_{22} is a polar amino acid residue, X_{23} is a polar charged amino acid residue, X_{24} is a polar amino acid residue, X_{25} is an aliphatic amino acid residue, X_{26} is a polar basic or neutral amino acid residue, X_{27} is a polar basic or neutral amino acid residue, and X_{28} is any amino acid residue. More specifically, X_{19} is asparagine, glutamine, or lysine; X_{20} is aspartic acid, glutamic acids, leucine, or
 10 isoleucine; X_{21} is valine, leucine, or isoleucine; X_{22} arginine, asparagine, or serine; X_{23} is glutamic acid or arginine; X_{24} is aspartic acid, glutamic acid, asparagine, glutamine, or arginine; X_{25} is valine, leucine, or isoleucine; X_{26} is arginine, asparagine, or lysine; X_{27} is asparagine or lysine; or X_{28} is glycine, alanine, or proline; or any combination of the foregoing.

15

In yet another embodiment, the peptide has the amino acid sequence

$DELAX_{29}X_{30}AQX_{31}WAX_{32}QCX_{33}X_{34}X_{35}X_{36}HD$ (SEQ ID NO:37), wherein X_{29} is any amino acid residue, X_{30} is an aliphatic amino acid residue, X_{31} is any amino acid residue, X_{32} is a polar neutral amino acid residue, X_{33} is a polar neutral or acidic amino acid

20 residue, X_{34} is an aromatic amino acid residue, X_{35} is an aliphatic amino acid residue or deleted, and X_{36} is any amino acid residue. More specifically, X_{29} is tyrosine, lysine, or histidine; X_{30} is isoleucine, leucine, or valine; X_{31} is threonine or valine; X_{32} is serine or asparagine; X_{33} is glutamine, asparagine, aspartic acid, or serine; X_{34} is phenylalanine to tyrosine; X_{35} is isoleucine, leucine, or deleted; or X_{36} is glycine, asparagine, or valine; or
 25 any combination of the foregoing.

In still another specific embodiment, the peptide has the amino acid sequence

$NX_{37}X_{38}X_{39}X_{40}X_{41}X_{42}X_{43}X_{44}BX_{45}FX_{46}KX_{47}GHYTOM$ (SEQ ID NO:38), wherein X_{37} is a

cyclic amino acid residue, X_{38} is a polar basic or neutral amino acid residue, X_{39} is any amino acid residue, X_{40} is any amino acid residue, X_{41} is a non-polar amino acid residue, X_{42} is any amino acid residue, X_{43} is a polar amino acid residue or glycine, X_{44} is a polar basic or neutral amino acid residue, X_{45} is a polar neutral amino acid or deleted, X_{46} is any amino acid residue, and X_{47} is a non-polar amino acid residue. More specifically, X_{37} is proline or tyrosine; X_{38} is asparagine, arginine, or histidine; X_{39} is lysine, threonine, or
 30

- valine; X₄₀ is aspartic acid, glycine, or lysine; X₄₁ is isoleucine, leucine, phenylalanine, or threonine; X₄₂ is glutamine, isoleucine, leucine, methionine, serine, or threonine; X₄₃ is asparagine, glutamic acid, histidine, lysine, or glycine, X₄₄ is aspartic acid, asparagine, glutamine, or serine; X₄₅ is asparagine or deleted, X₄₆ is alanine, arginine, leucine, isoleucine, or serine; or X₄₇ is isoleucine, leucine, threonine, or valine; or any combination of the foregoing.

In still another embodiment, the peptide has the amino acid sequence

- FX₄₆KX₄₇GHYTOMVWX₄₈X₄₉T (SEQ ID NO:39), wherein X₄₆ is any amino acid residue, X₄₇ is a non-polar amino acid residue, X₄₈ is a small side chain amino acid residue, and X₄₉ is a polar basic or neutral amino acid residue. More specifically, X₄₆ is alanine, arginine, leucine, isoleucine, or serine; X₄₇ is isoleucine, leucine, threonine, or valine; X₄₈ is glycine or alanine; or X₄₉ is lysine or asparagine; or any combination of the foregoing.

- 15 In another embodiment of the invention wherein two overlapping peptides are combined, the peptide has the amino acid sequence

- NX₃₇X₃₈X₃₉X₄₀X₄₁X₄₂X₄₃X₄₄BX₄₅FX₄₆KX₄₇GHYTOMVWX₄₈X₄₉T (SEQ ID NO:40), wherein X₃₇ is a cyclic amino acid residue, X₃₈ is a polar basic or neutral amino acid residue, X₃₉ is any amino acid residue, X₄₀ is any amino acid residue, X₄₁ is a non-polar amino acid residue, X₄₂ is any amino acid residue, X₄₃ is a polar amino acid residue or glycine, X₄₄ is a polar basic or neutral amino acid residue, X₄₅ is a polar neutral amino acid or deleted, X₄₆ is any amino acid residue, X₄₇ is a non-polar amino acid residue, X₄₈ is a small side chain amino acid residue, and X₄₉ is a polar basic or neutral amino acid residue. More specifically, X₃₇ is proline or tyrosine; X₃₈ is asparagine, arginine, or histidine; X₃₉ is lysine, threonine, or valine; X₄₀ is aspartic acid, glycine, or lysine; X₄₁ is isoleucine, leucine, phenylalanine, or threonine; X₄₂ is glutamine, isoleucine, leucine, methionine, serine, or threonine; X₄₃ is asparagine, glutamic acid, histidine, lysine, or glycine, X₄₄ is aspartic acid, asparagine, glutamine, or serine; X₄₅ is asparagine or deleted. X₄₆ is alanine, arginine, leucine, isoleucine, or serine; X₄₇ is isoleucine, leucine, threonine, or valine; X₄₈ is glycine or alanine; or X₄₉ is lysine or asparagine; or any combination of the foregoing.

In still another embodiment, the peptide has the amino acid sequence

X₅₀ZX₅₁X₅₂X₅₃X₅₄X₅₅HYLX₅₆CNYGPX₅₇GNX₅₈X₅₉X₆₀ (SEQ ID NO:40), wherein X₅₀ is a

non-polar amino acid, X₅₁ is a polar acidic or neutral amino acid residue, X₅₂ is a polar basic or neutral amino acid residue, X₅₃ is a non-polar amino acid residue, X₅₄ is a moderately polar amino acid residue, X₅₅ is a polar basic or neutral amino acid residue, X₅₆ is an aliphatic amino acid residue, X₅₇ is any amino acid residue, X₅₈ is any amino acid residue, X₅₉ is any amino acid residue, and X₆₀ is any amino acid residue. More specifically, X₅₀ is isoleucine, leucine, or methionine; X₅₁ is asparagine, aspartic acid, glutamine, or glutamic acid; X₅₂ is asparagine or lysine; X₅₃ is methionine or tryptophan; X₅₄ glutamine, histidine, or tyrosine; X₅₅ is asparagine, lysine, or threonine; X₅₆ isoleucine, leucine, or valine; X₅₇ is alanine, glycine, or serine; X₅₈ is aspartic acid, phenylalanine, or tyrosine; X₅₉ is glutamine, leucine, isoleucine, methionine, or phenylalanine; or X₆₀ is asparagine, aspartic acid, or glycine; or any combination of the foregoing.

In specific embodiments, the invention is directed to a peptide having an amino acid sequence selected from the group consisting of:

NNYCKIKCRKGIHTLCKFGT (SEQ ID NO: 8), or its homologue;
 GIHTLCKFGTSMKPNCGRNV (SEQ ID NO:9), or its homologue;
 NNYCKIKCRKGIHTLCKFGTGTSMKPNCGRNV (SEQ ID NO:42), or its homologue;
 KNEILKRHNDFRQNVAKGLE (SEQ ID NO:12), or its homologue;
 FRQNVAKGLETRGKPGPQPP (SEQ ID NO:13), or its homologue;
 KNEILKRHNDFRQNVAKGLETRGKPGPQPP (SEQ ID NO:43), or its homologue;
 DELAKIAQTWANQCDFNHDD (SEQ ID NO:16), or its homologue;
 NYKVGLQNSNFRKVGHYTQM (SEQ ID NO:22), or its homologue;
 FRKVGHYTQMVWGKT (SEQ ID NO:23), or its homologue;
 NYKVGLQNSNFRKVGHYTQMVWGKT (SEQ ID NO:44), or its homologue;
 and
 IEDNWyTHYLVCNYGPGGND (SEQ ID NO:26), or its homologue;

In specific embodiments, *infra*, peptides having SEQ ID NOS:8, 9, 12, 13, 16, 22, 23, and 26 are shown to be dominant in more than one strain of mouse.

In addition to homologous variants, allelic variant, consensus variant, and combined variant peptides of the invention, the present invention further provides a recombinant polypeptide comprising two or more peptides non-contiguously arranged relative to the native sequence of vespid venom antigen 5. Although the term recombinant generally refers to polypeptides generated by genetic engineering, which meaning is fully intended according to the present invention, the term recombinant is used herein more generally, to refer to polypeptides created by combination (or recombination) of non-contiguous immunodominant peptide fragment of vespid venom antigen 5. Such fragments may be from antigen 5 from the same species, antigen 5 peptides from different species, consensus antigen 5 peptides as described above, or any combination of the foregoing. In a specific embodiment, the peptides are selected from the group consisting of:

- NNYCKIKCRKGIHTLCKFGT (SEQ ID NO: 8), or its homologue;
- GIHTLCKFGTSMKPNCGRNV (SEQ ID NO:9), or its homologue;
- NNYCKIKCRKGIHTLCKFGTGTSMKPNCGRNV (SEQ ID NO:42), or its homologue;
- KNEILKRHNDFRQNVAKGLE (SEQ ID NO:12), or its homologue;
- FRQNVAKGLETRGKPGPQPP (SEQ ID NO:13), or its homologue;
- KNEILKRHNDFRQNVAKGLETRGKPGPQPP (SEQ ID NO:43), or its homologue;
- DELA KIAQTWANQCDFNHDD (SEQ ID NO:16), or its homologue;
- NYKVGLQNSNFRKVGHYTQM (SEQ ID NO:22), or its homologue;
- FRKVGHYTQMVWGKT (SEQ ID NO:23), or its homologue;
- NYKVGLQNSNFRKVGHYTQMVWGKT (SEQ ID NO:44), or its homologue;
- and
- IEDN WYTHYLVCNYGPGGND (SEQ ID NO:26), or its homologue.

In specific embodiments, the present invention is directed to the following homologous T cell peptides of hornet Ag5 from two species each of hornet (Dol a and Dol m), yellowjacket (Ves m and Ves v) and wasp (Pol a and Pol e) and one species of fire ant (Sol i) shown in Table 2. As noted above, there are two forms of Dol m 5.01 and 5.02, referred to as Dol m A and B here. Residues given below refer to Dol m 5.02 numbering, and gaps are added to obtain maximal sequence alignment. Where immunodominant peptides overlap, the fragment containing the overlapping sequences are provided, with the specific sequences demarcated by a line under the peptide group. In

such specific embodiments, the present invention contemplates both the individual immunodominant peptides and the larger peptide containing the overlapping segments.

Table 2. Homologues of Immunodominant Peptides

5

Peptides 1 and 2 (residue 1-20 and 11-30)

10

20

25

30

35

40

45

50

5

10

15

20

25

30

35

40

45

50

55

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235

240

245

250

255

260

265

270

275

280

285

290

295

300

305

310

315

320

325

330

335

340

345

350

355

360

365

370

375

380

385

390

395

400

405

410

415

420

425

430

435

440

445

450

455

460

465

470

475

480

485

490

495

500

505

510

515

520

525

530

535

540

545

550

555

560

565

570

575

580

585

590

595

600

605

610

615

620

625

630

635

640

645

650

655

660

665

670

675

680

685

690

695

700

705

710

715

720

725

730

735

740

745

750

755

760

765

770

775

780

785

790

795

800

805

810

815

820

825

830

835

840

845

850

855

860

865

870

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950

955

960

965

970

975

980

985

990

995

1000

1005

1010

1015

1020

1025

1030

1035

1040

1045

1050

1055

1060

1065

1070

1075

1080

1085

1090

1095

1100

1105

1110

1115

1120

1125

1130

1135

1140

1145

1150

1155

1160

1165

1170

1175

1180

1185

1190

1195

1200

1205

1210

1215

1220

1225

1230

1235

1240

1245

1250

1255

1260

1265

1270

1275

1280

1285

1290

1295

1300

1305

1310

1315

1320

1325

1330

1335

1340

1345

1350

1355

1360

1365

1370

1375

1380

1385

1390

1395

1400

1405

1410

1415

1420

1425

1430

1435

1440

1445

1450

1455

1460

1465

1470

1475

1480

1485

1490

1495

1500

1505

1510

1515

1520

1525

1530

1535

1540

1545

1550

1555

1560

1565

1570

1575

1580

1585

1590

1595

1600

1605

1610

1615

1620

1625

1630

1635

1640

1645

1650

1655

1660

1665

1670

1675

1680

1685

1690

1695

1700

1705

1710

1715

1720

1725

1730

1735

1740

1745

1750

1755

1760

1765

1770

1775

1780

1785

1790

1795

1800

1805

1810

1815

1820

1825

1830

1835

1840

1845

1850

1855

1860

1865

1870

1875

1880

1885

1890

1895

1900

1905

1910

1915

1920

1925

1930

1935

1940

1945

1950

1955

1960

1965

1970

1975

1980

1985

1990

1995

2000

2005

2010

2015

2020

2025

2030

2035

2040

2045

2050

2055

2060

2065

2070

2075

2080

2085

2090

2095

2100

2105

2110

2115

2120

2125

2130

2135

2140

2145

2150

2155

2160

2165

2170

2175

2180

2185

2190

2195

2200

2205

2210

2215

2220

2225

2230

2235

2240

2245

2250

2255

2260

2265

2270

2275

2280

2285

2290

2295

2300

2305

2310

2315

2320

2325

2330

2335

2340

2345

2350

2355

2360

2365

2370

2375

2380

2385

2390

2395

2400

2405

2410

2415

2420

2425

2430

2435

2440

2445

2450

2455

2460

2465

2470

2475

2480

2485

2490

2495

2500

2505

2510

2515

2520

2525

2530

2535

2540

2545

2550

2555

2560

2565

2570

2575

2580

2585

2590

2595

2600

2605

2610

2615

2620

2625

2630

2635

2640

2645

2650

2655

2660

2665

2670

2675

2680

2685

2690

2695

2700

2705

2710

2715

2720

2725

2730

2735

2740

2745

2750

2755

2760

2765

2770

2775

2780

2785

2790

2795

2800

2805

2810

2815

2820

2825

2830

2835

2840

2845

2850

2855

2860

2865

2870

2875

2880

2885

2890

2895

2900

2905

2910

2915

2920

2925

2930

2935

2940

2945

2950

2955

2960

2965

2970

2975

2980

2985

2990

2995

3000

3005

3010

3015

3020

3025

3030

3035

3040

3045

3050

3055

3060

3065

3070

3075

3080

3085

3090

3095

3100

3105

3110

3115

3120

3125

3130

3135

3140

3145

3150

3155

3160

3165

3170

3175

3180

3185

3190

3195

3200

3205

3210

3215

3220

3225

3230

3235

3240

3245

3250

3255

3260

3265

3270

3275

3280

3285

3290

3295

3300

3305

3310

3315

3320

3325

3330

3335

3340

3345

3350

3355

3360

3365

3370

3375

3380

3385

3390

3395

3400

3405

3410

3415

3420

3425

3430

3435

3440

3445

3450

3455

3460

3465

3470

3475

3480

3485

3490

3495

3500

3505

3510

3515

3520

3525

3530

3535

3540

3545

3550

3555

3560

3565

3570

3575

3580

3585

3590

3595

3600

3605

3610

3615

3620

3625

3630

3635

3640

3645

3650

3655

3660

3665

3670

3675

3680

3685

3690

3695

3700

3705

3710

3715

3720

3725

3730

3735

3740

3745

3750

3755

3760

3765

3770

3775

3780

3785

3790

3795

3800

3805

3810

3815

3820

3825

3830

3835

3840

3845

3850

3855

3860

3865

3870

3875

3880

3885

3890

3895

3900

3905

3910

3915

3920

3925

3930

3935

3940

3945

3950

3955

3960

3965

3970

3975

3980

3985

3990

3995

4000

4005

4010

4015

4020

4025

4030

4035

4040

4045

4050

4055

4060

4065

4070

4075

4080

4085

4090

4095

4100

4105

4110

4115

4120

4125

4130

4135

4140

4145

4150

4155

4160

4165

4170

4175

4180

4185

4190

4195

4200

4205

4210

4215

4220

4225

4230

4235

4240

4245

4250

4255

4260

4265

4270

4275

4280

4285

4290

4295

4300

4305

4310

4315

4320

4325

4330

4335

4340

4345

4350

4355

4360

4365

4370

4375

4380

4385

4390

4395

4400

4405

4410

4415

4420

4425

4430

4435

4440

4445

4450

4455

4460

4465

4470

4475

4480

4485

4490

4495

4500

4505

4510

4515

4520

4525

4530

4535

4540

4545

4550

4555

4560

4565

4570

4575

4580

4585

4590

4595

4600

4605

4610

4615

4620

4625

4630

4635

4640

4645

4650

4655

4660

4665

4670

4675

4680

4685

4690

4695

4700

4705

4710

4715

4720

4725

4730

4735

4740

4745

4750

4755

4760

4765

4770

4775

4780

4785

4790

4795

4800

4805

4810

4815

4820

4825

4830

4835

4840

4845

4850

4855

4860

4865

4870

4875

4880

4885

4890

4895

4900

4905

4910

4915

4920

4925

4930

4935

4940

4945

4950

4955

4960

4965

4970

4975

4980

4985

4990

4995

5000

5005

5010

5015

5020

5025

5030

5035

5040

5045

5050

5055

5060

5065

5070

5075

5080

5085

5090

5095

5100

5105

5110

5115

5120

5125

5130

5135

5140

5145

5150

5155

5160

5165

5170

5175

5180

5185

5190

5195

5200

5205

5210

5215

5220

5225

5230

5235

5240

5245

5250

5255

5260

5265

5270

5275

5280

5285

5290

5295

5300

5305

5310

5315

5320

5325

5330

5335

5340

5345

5350

5355

5360

5365

5370

5375

5380

5385

5390

5395

5400

5405

5410

5415

5420

5425

5430

5435

5440

5445

5450

5455

5460

5465

5470

5475

5480

5485

5490

5495

5500

5505

5510

5515

5520

5525

5530

5535

5540

5545

5550

5555

5560

5565

5570

5575

5580

5585

5590

5595

5600

5605

5610

5615

5620

5625

5630

5635

5640

5645

5650

5655

5660

5665

5670

5675

5680

5685

5690

5695

5700

5705

5710

5715

5720

5725

5730

5735

5740

5745

5750

5755

5760

5765

5770

5775

5780

5785

5790

5795

5800

5805

5810

5815

5820

5825

5830

5835

5840

5845

5850

5855

5860

5865

5870

5875

5880

5885

5890

5895

5900

5905

5910

5915

5920

5925

5930

5935

5940

5945

5950

5955

5960

5965

5970

5975

5980

5985

5990

5995

6000

6005

6010

6015

6020

6025

6030

6035

6040

6045

6050

6055

6060

6065

6070

6075

6080

6085

6090

6095

6100

6105

6110

6115

6120

6125

6130

6135

6140

6145

6150

6155

6160

6165

6170

6175

6180

6185

6190

6195

6200

6205

6210

6215

6220

6225

6230

6235

6240

6245

6250

6255

6260

6265

6270

6275

6280

6285

6290

6295

6300

6305

6310

6315

6320

6325

6330

6335

6340

6345

6350

6355

6360

6365

6370

6375

6380

6385

6390

6395

6400

6405

6410

6415

6420

6425

6430

6435

6440

6445

6450

6455

6460

6465

6470

6475

6480

6485

6490

6495

6500

6505

6510

6515

6520

6525

6530

6535

6540

6545

6550

6555

6560

6565

6570

6575

6580

6585

6590

6595

6600

6605

6610

6615

6620

6625

6630

6635

6640

6645

6650

6655

6660

6665

6670

6675

6680

6685

6690

6695

6700

6705

6710

6715

6720

6725

6730

6735

6740

6745

6750

6755

6760

6765

6770

6775

6780

6785

6790

6795

6800

6805

6810

6815

6820

6825

6830

6835

6840

6845

6850

6855

6860

6865

6870

6875

6880

6885

6890

6895

6900

6905

6910

6915

6920

6925

6930

6935

6940

6945

6950

69

Table 2 (cont.)

Peptide 15 and 20 (residue 141-160 and 151-165)					
5	Ves m	NPKKKFSeNn	FLKiGHYTMVWANT	SEQ ID NO:68	
	Ves v	NPKKKFSgNd	FLKtGHYTMVWANT	SEQ ID NO:69	
	Dol a	NPhKdlmhNN	FSKVGHYTMVWGKT	SEQ ID NO:70	
	Dol mA	NPKKGtigdnn	FSKVGHYTMVWGKT	SEQ ID NO:71	
	Dol mB	NYKvGlgnsN	FrKVGHYTMVWGKT	SEQ ID NO:44	
10	Pol a	NYNTGITKQN	FAKiGHYTMVWGKT	SEQ ID NO:72	
	Pol e	NYNTGITKQN	FAKiGHYTMVWGKT	SEQ ID NO:73	
	Sol i	NYNTGISFPSDDNILMKVEHYTQIVWAKT	SEQ ID NO:74		

15

20

Peptide 18 (residue 176-195)					
20	Ves m	IQEnWHKHLYLVCNYGPSGNF	SEQ ID NO:75		
	Ves v	IQEKWHKHLYLVCNYGPSGNF	SEQ ID NO:76		
	Dol a	IENKWHTHLYLVCNYGPAGNY	SEQ ID NO:77		
	Dol mA	IENNWHTHLYLVCNYGPAGNY	SEQ ID NO:78		
25	Dol mB	IEdNWyTHLYLVCNYGPgGnd	SEQ ID NO:26		
	Pol a	mENNMQNHYLICNYGPAGNY	SEQ ID NO:79		
	Pol e	iENkMQNHYLICNYGPAGNY	SEQ ID NO:80		
	Sol i	EPDNWTKHYLVCNYGPAGNV	SEQ ID NO:81		

30 The present invention further provides a pharmaceutical composition for treating vespid venom sensitivity, preferably where sensitivity to antigen 5 has been demonstrated, comprising any of the foregoing peptides of the invention and a pharmaceutically acceptable carrier. In a preferred aspect, the pharmaceutical composition comprises more than one peptide of the invention, thus greatly increasing the breadth of its effectiveness.

35

The invention naturally extends as well to a method for treating sensitivity to vespid venom comprising administering to a vespid venom allergic patient a therapeutically effective amount of a pharmaceutical composition of the invention. Preferably, the patient has been identified as sensitive to vespid venom antigen 5.

40

As demonstrated herein, the peptides of the invention are cross-reactive for a native testes protein. Thus, an important advance of the present invention is that it provides peptides specific for treatment of antigen 5-sensitive patients, regardless of the cause of this antigen 5 sensitivity. In other words, as demonstrated herein, whether a subject develops antigen 5 sensitivity from vespid stings, fire ant bites, or to testes protein, the peptides of the present invention provide an advantageous therapeutic agent for treating such sensitivity.

45

Thus, a primary object of the present invention is to provide immunodominant peptides from vespid venom antigen 5.

A corollary object is to provide such immunodominant peptides that produce T cell anergy
5 in subjects sensitive or allergic to vespid venom antigen 5.

Yet another object of the invention is to provide combinations of such immunodominant peptides for a broad treatment of vespid venom sensitive patients from a wide variety of MHC (HLA) backgrounds.

10

Still another object of the invention is to provide consensus immunodominant T cell epitopes of vespid venom antigen 5s.

These and other objects of the present invention can be better appreciated by reference to
15 the following drawings, the Detailed Description of the Invention, and the Examples.

ABBREVIATIONS

	Ag5, Ag 5	antigen 5
	<i>Dol m Dolichovespula maculata</i>	white face hornet
20	<i>Dol a D. arenaria</i>	yellow hornet
	<i>Pol a Polistes annularis</i>	wasp
	<i>Pol e P. exclamans</i>	wasp
	<i>Ves m Vespula maculifrons</i>	yellowjacket
	<i>Ves v V. vulgaris</i>	yellowjacket
25	n-, r-	natural, recombinant (respectively)
	PCR	polymerase chain reaction
	TCR	T cell receptor for antigen
	tpx	testes protein

30

BRIEF DESCRIPTION OF THE DRAWINGS

FIGURE 1. Amino acid sequence of vespid antigen 5s. Antigen 5s were isolated from yellowjackets (*Vespula maculifrons* and *vulgaris*), hornets (*Dolichovespula arenaria* and *maculata*) and wasps (*Polistes annularis* and *exclamans*). The sequences given are in the

order of *Ves m V*, *Ves v V*, *Dol à V*, *Dol m VA* and *VB*, *Pol a V*, and *Pol e V* respectively. Residues which are common to all sequences are given on separate lines beneath the sequences; residues underlined or dotted represent conserved or variable regions, respectively.

5

FIGURE 2. Peptide fragments of *Dol m* Antigen 5.2.

FIGURE 3. Sequence similarity of hornet Ag5 and mouse and human tpx's. **Bold characters** indicate those residues of human and mouse tpx's which are identical with those of Ag5. The underlines regions of greater than 7 residues in length have greater than 67% sequence identity, and each is interspersed with fewer than 4 substituted residues.

FIGURE 4. Stimulation index profile of BALB/c mice spleens after 4 immunizations with natural (n)-Ag5. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant antigen 5 control were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 2200 cpm.

FIGURE 5. Stimulation index profile of BALB/c mice spleens after 4 immunizations with recombinant (r)-Ag5. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 4700 cpm.

25

FIGURE 6. Stimulation index profile of BALB/c mice spleens after 4 immunizations with r-Ag5. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 3970 cpm.

30

FIGURE 7. Stimulation index profile of BALB/c mice spleens after 4 immunizations with r-frag IN (residue 1-114 of antigen 5). *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill"

hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 4690 cpm.

5 FIGURE 8. Stimulation index profile of BALB/c mice spleens after 4 immunizations with r-frag C (residue 151-204 of antigen 5). *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was
10 6620 cpm.

FIGURE 9. Stimulation index profile of BALB/c mice spleens after 5 immunizations with r-frag IN. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M
15 ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 4410 cpm.

FIGURE 10. Stimulation index profile of BALB/c mice spleens after 5 immunizations with r-frag C. *In vitro* proliferation assays were performed with peptides
20 at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 7520 cpm.

FIGURE 11. Stimulation index profile of ASW/sn mice spleens after 5 immunizations with r-Ag5. *In vitro* proliferation assays were performed with peptides at
25 three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 5640 cpm.

30 FIGURE 12. Stimulation index profile of P/J mice spleens after 5 immunizations with r-Ag5. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 1730 cpm.

FIGURE 13. Stimulation index profile of C3H/He mice spleens after 5 immunizations with r-Ag5. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 3130 cpm.

FIGURE 14. Stimulation index profile of C57Bl/6 mice spleens after 5 immunizations with r-Ag5. *In vitro* proliferation assays were performed with peptides at three concentrations: 1×10^{-5} M (open bar), 1.25×10^{-6} M ("uphill" hatch), and 1.56×10^{-7} M ("downhill" hatch). The corresponding concentrations of recombinant Antigen 5 were 1.0×10^{-6} M, 1.25×10^{-7} , and 1.56×10^{-8} M. The blank was 1050 cpm.

FIGURE 15. Stimulation index profile of BALB/c mice spleen cells after 4 immunizations with Tpx-N. Stimulation was tested with peptide or protein at three different concentrations: Open bars -- 1.0×10^{-5} M peptide or Frag IN, 2.5×10^{-6} M Tpx-N; "uphill" hatch -- 1.25×10^{-6} M peptide or Frag IN, 3.13×10^{-7} M, Tpx-N; "downhill" hatch -- 1.5×10^{-7} M peptide and Frag IN, 3.91×10^{-8} M Tpx-N; and stippled -- 1.96×10^{-8} M Frag IN, 4.89×10^{-9} M Tpx-N. The blank was 7270 cpm.

FIGURE 16. Stimulation index profile of BALB/c mice spleen cells after 4 immunizations with Tpx-C. Stimulation was tested with peptide or protein at three different concentrations: Open bars - 1×10^{-5} M peptide, 5×10^{-6} M Frag C or Tpx-C; "uphill" hatch -- 1.25×10^{-6} M peptide; 6.25×10^{-7} M Frag C or Tpx - C; "downhill" hatch -- 1.56×10^{-7} M peptide, 7.8×10^{-8} M Frag C, Tpx - C; and stippled -- 9.75×10^{-9} M Frag - C or Tpx - C. The blank was 5110 cpm.

DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention provides the sequences of immunodominant peptides of vespid venom antigen 5, and corresponding peptides from other antigens, such as fire ant Sol i 3.

The term "immunodominant peptide" is used herein to refer to a peptide that contains a T cell epitope of the antigen, such that T cells immunized with the antigen will be stimulated

when contacted with the peptide. The term T cell epitope is used herein in conformity with the definition in immunology. In one embodiment, an immunodominant epitope is an epitope that induces a greater than about 4-fold increase in stimulation of immunized T cells in an *in vitro* stimulation assay, particularly a greater than 2-fold, preferably greater than 4-fold, and more preferably greater than 6-fold increase in the stimulation index in a ³H-thymidine incorporation assay. In another embodiment, an immunodominant peptide is a peptide that induces proliferation or stimulation of different MHC-restricted populations of immunized T cells.

- 10 An immunodominant peptide of the invention may contain one epitope, or it may contain more than one epitope.

An immunomodulatory peptide" is a peptide that affects immune response *in vivo*. Preferably, an immunomodulatory peptide of the invention induces T cell anergy when
15 administered to a subject.

- The invention also provides pharmaceutical compositions effective for the treatment of a vespid venom allergen-specific allergic condition comprising a polypeptide of the invention, and methods for treating such allergic conditions comprising administering a
20 therapeutically effective dose of the pharmaceutical compositions of the invention.

The polypeptides of the invention can also be useful for diagnosis of vespid venom-specific allergic conditions.

- 25 As used herein, the term "vespid venom allergen" refers to a protein found in the venom of a vespid, to which susceptible people are sensitized on exposure to the sting of the insect. While most antigens are characterized by being reactive with specific IgG class antibodies, an allergen is characterized by also being reactive with IgE type antibodies. The IgE type antibodies are responsible for mediating the symptoms of an allergic
30 condition, *i.e.*, immediate-type hypersensitivity. According to the invention, the vespid venom allergen of interest is vespid venom antigen 5.

As herein, the term "vespid" is used according to the practice of those in the field of allergy, and refers to insects belonging to the worldwide family of Vespidae, *i.e.*, social

wasps including hornets, yellowjackets, and paper wasps. In particular, vespids include the subfamilies Vespinae and Polistinae. More particularly, the vespids include the genera *Vespa* Linnaeus, *Vespula* Thomson, *Dolichovespula* Rohwer, and *Polistes* Latreille. Species in the genus *Vespula* include but are not limited to *V. germanica* (Fab.), *V.*
 5 *squamosa* (Drury), *V. maculifrons* (Buysson), *V. flavopilosa* (Jacobson), *V. vulgaris* (L.), and *V. pennsylvanica* (Saussure). Species in the genus *Polistes* include but are not limited to *P. annularis* (Linnaeus), *P. exclamans* (Viereck), *P. metricus* (Say), *P. fuscatus* (Fabricius), and *P. apachus* (Saussure). Species in the genus *Dolichovespula* include but are not limited to *D. maculata* (L.) and *D. arenaria* (Fab.). Species in the genus *Vespa*
 10 include but are not limited to *V. crabro* (L.) and *V. orientalis* (Linnaeus).

As used herein, the term "immunomodulatory" refers to an ability to increase or decrease an antigen-specific immune response, either at the B cell or T cell level. Immunomodulatory activity can be detected *e.g.*, in T cell proliferation assays, by
 15 measurement of antibody production, lymphokine production or T cell responsiveness. In particular, in addition to affects on T cell responses, the immunomodulatory polypeptides of the invention may bind to immunoglobulin (*i.e.*, antibody) molecules on the surface of B cells, and affect B cell responses as well.

20 For the sake of clarity, the present invention is described in detail in sections relating to an immunomodulatory fragment of a vespid venom antigen 5, or derivatives and analogs of the vespid venom antigen 5, assays with the immunomodulatory vespid venom antigen 5, fragments, or derivatives or analogs thereof, and finally therapeutic and diagnostic uses of the vespid venom antigen 5 fragments, or derivatives or analogs thereof.

25

Immunodominant Peptides of Vespid Venom Ag5

An immunodominant peptide of the invention is a peptide comprising at least one T cell epitope, with a minimum size of approximately 8 amino acid residues (the minimum length generally regarded as requisite for antigen-specific recognition by a T cell). A
 30 peptide comprising at least one T cell epitope is capable of eliciting a T cell response, such as stimulation or T cell anergy (tolerization).

The immunodominant peptides of the invention have been identified in murine backgrounds. However, those peptides that are antigenic (comprise a T cell epitope) in

more than one mouse strain are excellent candidates for immunodominant epitopes in antigen 5 sensitive humans. As pointed out above, on the basis of findings with Fel d 1 and Der p 1, it is reasonable to conclude that the T cell epitope data of a vespid venom antigen 5 found in mice, particularly in multiple strains, will be applicable for humans.

5

Also contemplated by the term "immunodominant peptide" are peptides which join two or more discontinuous epitopes (or the specific peptides disclosed herein) in a single polypeptide. In one aspect, such "aggregate" polypeptides can be prepared chemically, *e.g.*, by crosslinking peptides, or using condensation techniques to form peptide bonds.

10 Alternatively, synthetic DNA molecules encoding such aggregate peptides can be prepared, inserted in a recombinant expression vector, and the aggregate peptide expressed by host cells transformed or transfected with the expression vector and cultured under conditions that allow for expression of the aggregate polypeptide.

15 Immunodominant peptides of the invention can also be prepared as fusion proteins.

Amino acids used for peptide synthesis may be standard Boc (N^{α} -amino protected N^{α} -t-butyloxycarbonyl) amino acid resin with the standard deprotecting, neutralization, coupling and wash protocols of the original solid phase procedure of Merrifield [1963, *J. Am.*
20 *Chem. Soc.* **85**:2149-2154], or the base-labile N^{α} -amino protected 9-fluorenylmethoxycarbonyl (Fmoc) amino acids first described by Carpino and Han [1972, *J. Org. Chem.* **37**:3403-3409]. Both Fmoc and Boc N^{α} -amino protected amino acids can be obtained from Fluka, Bachem, Advanced Chemtech, Sigma, Cambridge Research Biochemical, Bachem, or Peninsula Labs or other chemical companies familiar
25 to those who practice this art. In addition, the method of the invention can be used with other N^{α} -protecting groups that are familiar to those skilled in this art. Solid phase peptide synthesis may be accomplished by techniques familiar to those in the art and provided, for example, in Stewart and Young, 1984, *Solid Phase Synthesis*, Second Edition, Pierce Chemical Co., Rockford, IL; Fields and Noble, 1990, *Int. J. Pept. Protein*
30 *Res.* **35**:161-214, or using automated synthesizers, such as sold by ABS.

Alternatively, immunodominant peptides of the invention can be prepared by recombinant DNA techniques. Thus, in accordance with the present invention there may be employed conventional molecular biology, microbiology, and recombinant DNA techniques within

the skill of the art. Such techniques are explained fully in the literature [see, *e.g.*, Sambrook, Fritsch & Maniatis, *Molecular Cloning: A Laboratory Manual*, Second Edition (1989) Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York [herein "Sambrook et al., 1989"]; *DNA Cloning: A Practical Approach*, Volumes I and II [D.N. Glover ed. 1985]; *Oligonucleotide Synthesis* [M.J. Gait ed. 1984]; *Nucleic Acid Hybridization* [B.D. Hames & S.J. Higgins eds. (1985)]; *Transcription And Translation* [B.D. Hames & S.J. Higgins, eds. (1984)]; *Animal Cell Culture* [R.I. Freshney, ed. (1986)]; *Immobilized Cells And Enzymes* [IRL Press, (1986)]; B. Perbal, *A Practical Guide To Molecular Cloning* (1984); F.M. Ausubel et al. (eds.), *Current Protocols in Molecular Biology*, John Wiley & Sons, Inc. (1994)].

Peptides comprising at least two regions, each region comprising at least one T cell epitope of an antigen 5 are also within the scope of the invention. Each region of such peptides is derived from the same or from different antigen 5. Isolated peptides, each comprising at least two T cell epitopes of antigen 5, are particularly desirable for increased therapeutic effectiveness. Peptides immunologically related by T cell cross-reactivity, which are capable of reacting with the same T cells as a peptide of the invention, are also contemplated.

As noted above, isolated peptides of the invention can be produced by recombinant DNA techniques in a host cell transformed with a nucleic acid having a sequence encoding such peptide. The isolated peptides of the invention can also be produced by chemical synthesis. In certain limited situations, isolated peptides can be produced by chemical cleavage of the protein allergen. When a peptide is produced by recombinant techniques, host cells transformed with a nucleic acid having a sequence encoding the peptide or the functional equivalent of the nucleic acid sequence are cultured in a medium suitable for the cells and peptides can be purified from cell culture medium, host cells, or both using techniques known in the art for purifying peptides and proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, or immunopurification with antibodies specific for the peptide, antigen 5 from which the peptide is derived, or a portion thereof. Isolated peptides of the invention are substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or substantially free of chemical precursors or other chemicals when synthesized chemically.

Naturally, the present invention provides expression vectors and host cells transformed to express the peptides. A nucleic acid sequence coding for a peptide of the invention may be expressed in bacterial cells such as *E. coli*, insect cells (baculovirus), yeast, or mammalian cells such as Chinese hamster ovary cells (CHO). Suitable expression
5 vectors, promoters, enhancers, and other expression control elements may be found in Sambrook et al. *Molecular Cloning: A Laboratory Manual*, second edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1989). Other suitable expression vectors, promoters, enhancers, and other expression elements are known to those skilled in the art. Many yeast and bacterial vectors are commercially available or on
10 deposit at various depositories. Baculovirus and mammalian expression systems are also available.

For expression in *E. coli*, suitable expression vectors include, among others, pTRC [Amann et al., *Gene*, **69**:301-315, (1988)], pGEX (Amrad Corp., Melbourne, Australia);
15 pMAL (N.E. Biolabs, Beverly, MA); pRIT5 (Pharmacia, Piscataway, NJ); pET-11d (Novagen, Madison, WI) [Jameel et al., *J. Virol.* **64**:3963-3966 (1990)]; pQE12 (QIAGEN, Chatsworth, CA); and pSEM [Knapp et al., *BioTechniques*, **8**:280-281 (1990)]. The use of pTRC, and pET-11d, for example, will lead to the expression of unfused protein. The use of pMAL, pRIT5, pSEM, pQE12, and pGEX will lead to the
20 expression of allergen fused to maltose E binding protein (pMAL), protein A (pRIT5), truncated β -galactosidase (PSEM), hexahistidine (pQE12), or glutathione S-transferase (pGEX). When a polypeptide allergen is expressed as a fusion protein, it is particularly advantageous to introduce an enzymatic cleavage site at the fusion junction between the carrier protein and a polypeptide. The immunodominant polypeptide (harboring the
25 peptide of the invention) may then be recovered from the fusion protein through enzymatic cleavage at the enzymatic site and biochemical purification using conventional techniques for purification of proteins and peptides. Suitable enzymatic cleavage sites include those for blood clotting Factor Xa or thrombin for which the appropriate enzymes and protocols for cleavage are commercially available from, for example, Sigma Chemical Company,
30 St. Louis, MO and N.E. Biolabs, Beverly, MA. The different vectors also have different promoter regions allowing constitutive or inducible expression with, for example, IPTG induction (PRTC [Amann et al., (1988), *supra*]; pET-11d, Novagen, Madison, WI) or temperature induction (pRIT5, Pharmacia, Piscataway, NJ).

Human T cell stimulating activity can be tested by culturing T cells obtained from an individual sensitive to vespid venom (*i.e.*, an individual who has an IgE mediated immune response to antigen 5) with an immunodominant peptide of the invention and determining whether proliferation of T cells occurs in response to the peptide as measured, *e.g.*, by cellular uptake of tritiated thymidine. Stimulation indices for responses by T cells to peptides can be calculated as the maximum CPM in response to a peptide divided by the control CPM. A T cell stimulation index (S.I.) equal to or greater than two times the background level is considered "positive." Positive results are used to calculate the mean stimulation index for each peptide for the group of peptides tested. Peptides of this invention comprise at least one T cell epitope and have a mean T cell stimulation index of greater than or equal to 2.0. Preferably the S.I. value is greater than four. A peptide having a T cell stimulation index of greater than or equal to 2.0, and more preferably greater than or equal to 4.0, is considered useful as a therapeutic agent. Preferred peptides have a mean T cell stimulation index of at least 2.5, more preferably at least 3.5, even more preferably at least 4.0, and most preferably at least 5.0.

In addition, preferred peptides have a positivity index (P.I.) of at least about 100, more preferably at least 150, even more preferably at least about 200 and most preferably at least about 250. The positivity index for a peptide is determined by multiplying the mean T cell stimulation index by the percent of individuals, in a population of individuals sensitive to vespid venom (*e.g.*, preferably at least 9 individuals, more preferably at least 16 individuals or more, more preferably at least 29 individuals or more, or even more preferably at least 30 individuals or more), who have T cells that respond to the peptide. Thus, the positivity index represents both the strength of a T cell response to a peptide (S.I.) and the frequency of a T cell response to a peptide in a population of individuals sensitive to vespid venom.

In order to determine precise T cell epitopes by, for example, fine mapping techniques, a peptide having T cell stimulating activity and thus comprising at least one T cell epitope as determined by T cell biology techniques is modified by addition or deletion of amino acid residues at either the amino or carboxyl terminus of the peptide and tested to determine a change in T cell reactivity to the modified peptide. If two or more peptides which share an area of overlap in the native protein sequence are found to have T cell stimulating activity, preferably human T cell stimulating activity, as determined by T cell biology

techniques, additional peptides can be produced comprising all or a portion of such peptides and these additional peptides can be tested by a similar procedure. Following this technique, peptides are selected and produced recombinantly or synthetically.

Peptides are selected based on various factors, including the strength of the T cell

- 5 response to the peptide (*e.g.*, stimulation index), the frequency of the T cell response to the peptide in a population of individuals sensitive to vespid venom, and the potential cross-reactivity of the peptide with other antigen 5 homologues. The physical and chemical properties of these selected peptides (*e.g.*, solubility, stability) are examined to determine whether the peptides are suitable for use in therapeutic compositions or whether
- 10 the peptides require modification as described herein. The ability of the selected peptides or selected modified peptides to stimulate human T cells (*e.g.*, induce proliferation, lymphokine secretion) is determined.

- Additionally, preferred peptides of the invention do not bind immunoglobulin E (IgE), or
- 15 bind IgE to a substantially lesser extent than the protein allergen from which the peptide is derived binds IgE. The major complications of standard immunotherapy are IgE-mediated response such as anaphylaxis. Immunoglobulin E is a mediator of anaphylactic reactions which result from the binding and cross-linking of antigen to IgE on mast cells or basophils and the release of mediators (*e.g.*, histamine, serotonin, eosinophil chemotactic
- 20 factors). Thus, anaphylaxis in a substantial percentage of a population of individuals sensitive to vespid venom could be avoided by the use in immunotherapy of a peptide of peptides which do not bind IgE in a substantial percentage (*e.g.*, at least about 75 %) of a population of individuals sensitive to vespid venom allergen, or if the peptide binds IgE, such binding does not result in the release of mediators from mast cells or basophile. The
- 25 risk of anaphylaxis could be reduced by the use in immunotherapy of a peptide or peptides which have reduced IgE binding. Thus, peptides which have minimal IgE stimulating activity are desirable for therapeutic effectiveness. Minimal IgE stimulating activity refers to IgE production that is less than the amount of IgE production and/or IL-4 production stimulated by the native protein allergen.

30

A peptide of the invention, when administered to a vespid venom-sensitive individual, is capable of modifying the allergic response of the individual to the allergen. Particularly, peptides of the invention comprising at least one T cell epitope of antigen 5 when administered to an individual sensitive to vespid venom are capable of modifying T cell

response of the individual to the allergen. As used herein, modification of the allergic response of an individual to vespid venom antigen 5 can be defined as non-responsiveness or diminution in symptoms upon exposure to antigen 5, as determined by standard clinical procedures [see *e.g.*, Varney et al., *British Medical Journal*, **302**:265-169 (1990)],
5 including diminution in antigen 5 induced asthmatic symptoms. As referred to herein, a diminution in symptoms includes any reduction in allergic response of an individual to the allergen after the individual has completed a treatment regimen with a peptide or protein of the invention. This diminution may be subjective (*i.e.*, the patient feels more comfortable in the presence of the allergen). Diminution in symptoms can be determined
10 clinically as well, using standard skin tests as is known in the art.

As a result of the work described herein, peptides derived from antigen 5 comprising at least one T cell epitope can be produced. T cell epitopes are believed to be involved in initiation and perpetuation of the immune responses to allergen(s), which are responsible
15 for the clinical symptoms of vespid venom. These T cell epitopes are thought to trigger early events at the level of the T helper cell by binding to an appropriate HLA (MHC) molecule on the surface of an antigen presenting cell and stimulating the relevant T cell subpopulation. These events lead to T cell proliferation, lymphokine secretion, local inflammatory reactions, the recruitment of additional immune cells to the site, and
20 activation of the B cell cascade leading to production of antibodies. One isotype of these antibodies, IgE, is fundamentally important in the development of allergic symptoms and its production is influenced early in the cascade of events, at the level of the T helper cell, by the nature of the lymphokines secreted. A T cell epitope is the basic element or smallest unit of recognition by a T cell receptor where the epitope comprises amino acid
25 residues essential to receptor recognition. Amino acid sequences which mimic those of T cell epitopes and which modify the allergic response to vespid venom antigen 5 are within the scope of this invention.

Exposure of vespid venom allergic patients to peptides of the present invention may
30 tolerize or anergize appropriate T cell subpopulations such that they become unresponsive to vespid venom allergen(s), *e.g.*, antigen 5, and do not participate in mounting an immune response upon such exposure. In addition, administration of a peptide of the present invention may modify the lymphokine secretion profile as compared with exposure to the naturally-occurring vespid venom allergin or portion thereof (*e.g.*, result in a

- decrease of IL-4 and/or an increase in IL-2). Furthermore, exposure to a peptide of the invention may influence T cell subpopulations which normally participate in a response to vespid venom allergen(s) such that these T cells are drawn away from the site(s) of normal exposure to the allergen (*e.g.*, nasal mucosa, skin, and lung) towards the site(s) of therapeutic administration of the peptide. This redistribution of T cell subpopulations may ameliorate or reduce the ability of an individual's immune system to mount an immune response at the site of normal exposure to the antigen 4, resulting in a diminution in allergic symptoms.
- 10 Isolated peptides of the invention comprise at least one T cell epitope of vespid venom antigen 5 and accordingly, the peptide comprises at least approximately eight amino acid residues of the protein allergen. For purposes of therapeutic effectiveness, therapeutic compositions of the invention preferably comprise at least two T cell epitopes of antigen 5. Accordingly, isolated peptides of the invention preferably comprise at least two T cell
- 15 epitopes and accordingly, the peptide comprises at least approximately nine amino acid residues, and preferably at least 15 amino acid residues. Additionally, therapeutic compositions of the invention preferably comprise a sufficient percentage of the T cell epitopes of the entire protein allergen such that a therapeutic regimen of administration of the composition to an individual sensitive to vespid venom results in T cells of the
- 20 individual being tolerized to the protein allergen. Synthetically produced peptides of the invention comprising up to approximately 35 amino acid residues in length, and more preferably up to approximately 20 amino acid residues in length are particularly desirable. Increases in length beyond this point may result in difficulty in peptide synthesis as well as retention of an undesirable property (*e.g.*, immunoglobulin binding or enzymatic activity)
- 25 due to maintenance of conformational similarity between the peptide and the protein allergen from which it is derived.

Another embodiment of the present invention provides peptides comprising at least two regions, each region comprising at least one T cell epitope of vespid venom antigen 5 and

30 accordingly, each region comprises at least approximately seven amino acid residues. These peptides comprising at least two regions can comprise as many amino acid residues as desired and preferably comprise at least about 14, even more preferably about 25, and most preferably at most about 35 amino acid residues of antigen 5. Each region of such peptide preferably comprises up to 20 amino acid residues in length, more preferably up

to 15 residues in length and most preferably up to 10 amino acid residues in length as increases in length of a region may result in difficulty in peptide synthesis as well as retention of an undesirable property (*e.g.*, immunoglobulin binding or enzymatic activity) due to maintenance of conformational similarity between the peptide and the protein

- 5 allergen from which it is derived. If desired, the amino acid sequences of the regions can be produced and joined by a linker to increase sensitivity to processing by antigen-presenting cells. Such a linker can be any non-epitope amino acid sequence or other appropriate linking or joining agent. To obtain preferred peptides comprising at least two regions, each comprising at least one T cell epitope, the regions are arranged in a
- 10 configuration different from a naturally-occurring configuration of the regions in the allergen or a combination of different antigen 5s. For example, the regions containing T cell epitope(s) can be arranged in a contiguous configuration and can preferably be derived from the same antigen or a combination of antigen 5 homologues from different species. Noncontiguous is defined as an arrangement of regions containing T cell epitope(s) which
- 15 is different from that of an amino acid sequence present in the protein allergen from which the regions are derived. Furthermore, the noncontiguous regions containing T cell epitopes can be arranged in a nonsequential order (*e.g.*, in an arrangement different from the order of the arrangement found in the native protein allergen from which the region containing T cell epitope(s) are derived).

20

The present invention further provides for modification or derivatization of peptides in a library. Modifications of peptides are well known to one of ordinary skill, and include phosphorylation, carboxymethylation, and acylation. Modifications may be effected by chemical or enzymatic means.

25

- It is also possible to modify the structure of a peptide of the invention for such purposes as increasing solubility, enhancing therapeutic or preventive efficacy, or stability (*e.g.*, shelf life *ex vivo*, and resistance to proteolytic degradation *in vivo*). A modified peptide can be produced in which the amino acid sequence has been altered, such as by amino
- 30 acid substitution, deletion, or addition, to modify immunogenicity and/or reduce allergenicity, or to which a component has been added for the same purpose.

For example, a peptide can be modified so that it maintains the ability to induce T cell anergy and bind MHC proteins without the ability to induce a strong proliferative response

or possibly any proliferative response when administered in immunogenic form. In this instance, critical binding residues for the T cell receptor can be determined using known techniques (*e.g.*, substitution of each residue and determination of the presence or absence of T cell reactivity). Those residues shown to be essential to interact with the T cell
5 receptor can be modified by replacing the essential amino acid with another, preferably similar amino acid residue (a conservative substitution) whose presence is shown to enhance, diminish but not eliminate, or not affect T cell reactivity. In addition, those amino acid residues that are not essential for T cell receptor interaction can be modified by being replaced by another amino acid whose incorporation may enhance, diminish or
10 not affect T cell reactivity, but does not eliminate binding to relevant MHC.

Additionally, peptides of the invention can be modified by replacing an amino acid shown to be essential to interact with the MHC protein complex with another, preferably similar amino acid residue (conservative substitution) whose presence is shown to enhance,
15 diminish but not eliminate, or not effect T cell activity. In addition, amino acid residues which are not essential for interaction with the MHC protein complex but which still bind the MHC protein complex can be modified by being replaced by another amino acid whose incorporation may enhance, not effect, or diminish but not eliminate T cell reactivity. Preferred amino acid substitutions for non-essential amino acids include, but
20 are not limited to, substitutions with alanine, glutamic acid, or a methyl amino acid.

Another example of a modification of peptides is substitution of cysteine residues preferably with serine, threonine, leucine or glutamic acid to minimize dimerization via disulfide linkages.

25

In order to enhance stability and/or reactivity, peptides can also be modified to incorporate one or more polymorphism in the amino acid sequence of a protein allergen resulting from natural allelic variation or from homologous peptides. Additionally, D-amino acids, non-natural amino acids, or non-amino acid analogs can be substituted or added to produce a
30 modified peptide within the scope of this invention. Furthermore, peptides of the present invention can be modified using the polyethylene glycol (PEG) method of A. Schon and co-workers [Wie et al., *supra*] to produce a peptide conjugated with PEG. In addition, PEG can be added during chemical synthesis of a peptide of the invention. Modifications of peptides or portions thereof can also include reduction/alkylation [Tarr in: *Methods of*

Protein Microcharacterization, J.E. Silver ed., Humana Press: Clifton, NJ, pp. 155-194) (1986)]; acylation [Tarr, *supra*]; esterification [Tarr, *supra*]; chemical coupling to an appropriate carrier [Mishell and Shiigi, eds., *Selected Methods in Cellular Immunology*, WH Freeman: San Francisco, CA (1980); U.S. Patent 4,939,239]; or mild formalin
5 treatment [Marsh, *Int. Arch. Allergy App. Immunol.*, **42**:199-215 (1971)].

To facilitate purification and potentially increase solubility of peptides of the invention, it is possible to add reporter group(s) to the peptide backbone. For example, polyhistidine can be added to a peptide to purify the peptide on immobilized metal ion affinity
10 chromatography [Hochuli, E. et al., *Bio/Technology* **6**:1321-1325 (1988)]. In addition, specific endoprotease cleavage sites can be introduced, if desired, between a reporter group and amino acid sequences of a peptide to facilitate isolation of peptides free of irrelevant sequences. In order to successfully desensitize an individual to a protein antigen, it may be necessary to increase the solubility of a peptide by adding functional
15 groups to the peptide or by not including hydrophobic T cell epitopes or regions containing hydrophobic epitopes in the peptides.

To potentially aid proper antigen processing of T cell epitopes within a peptide, canonical protease sensitive sites can be recombinantly or synthetically engineered between regions,
20 each comprising at least one T cell epitope. For example, charged amino acid pairs, such as KK or RR, can be introduced between regions within a peptide during recombinant construction of the peptide. The resulting peptide can be rendered sensitive to proteolytic cleavage, *e.g.*, with cathepsin and/or other trypsin-like enzymes, to generate portions of the peptide containing one or more T cell epitopes. In addition, such charged amino acid
25 residues can result in an increase in solubility of a peptide.

Fatty acyl peptide derivatives may also be prepared. For example, and not by way of limitation, a free amino group (N-terminal or lysyl) may be acylated, *e.g.*, myristoylated. In another embodiment an amino acid comprising an aliphatic side chain of the structure -
30 $(CH_2)_nCH_3$ may be incorporated in peptides of the library. This and other peptide-fatty acid conjugates suitable for use in the present invention are disclosed in U.K. Patent GB-8809162.4, International Patent Application PCT/AU89/00166.

Activity Assays With Peptides of the Invention

Numerous assays are known in immunology for evaluating the immunomodulatory activity of an antigen. For example, the immunodominant vespid venom can be used in diagnostic assays for allergic diseases, which are described in detail, *infra*. In general, such peptides
5 can be tested for the ability to induce proliferation of T cells from allergic subjects, without reacting with antibodies specific for antigen 5. Preferably, such antibodies that are not reactive with the peptides of the invention in the diagnostic assay are of the IgE class. It is important to note that natural allergen-specific antibodies have been found to bind weakly to denatured vespid venom allergens.

10

The peptides of the invention can be tested in a proliferation assay for T cell responses. Generally, lymphocytes from a sensitized host are obtained. The host can be a mouse that has been immunized with a vespid venom antigen 5, or with a crossreactive protein, such as testes specific protein. Using techniques that are well known in the art, T lymphocyte
15 response to the peptide can be measured *in vitro*. In a specific embodiment, *infra*, T cell responses are detected by measuring incorporation of ³H-thymidine, which increases with DNA synthesis associated with proliferation. Cell proliferation can also be detected using an MTT assay [Mossman, *J. Immunol. Methods*, **65**:55-63 (1983); Niks and Otto, *J. Immunol. Methods*, **130**:140-151 (1990)]. Alternatively, lymphokine production assays
20 can be practiced according to the present invention. In one embodiment, lymphokine production can be assayed using immunological or co-stimulation assays [see, *e.g.*, Fehlner et al., *J. Immunol.*, **146**:799 (1991)] or using the ELISPOT technique [Czerkinsky, et al., *J. Immunol. Methods*, **110**:29 (1988)]. Alternatively, mRNA for lymphokines can be detected, *e.g.*, by amplification [see Brenner, et al., *Biotechniques*,
25 **7**:1096 (1989)] or *in situ* hybridization [see, *e.g.*, Kasaian and Biron, *J. Immunol.*, **142**:1287 (1989)]. Of particular interest are those individuals whose T cells produce lymphokines associated with IgE isotype switch events, *e.g.*, IL-4 and IL-5 [Purkeson and Isakson, *J. Exp. Med.*, **175**:973-982 (1992)]. Also of interest are the peptide fragments of the vespid venom antigen 5 that contain epitopes recognized by T cells involved in IgE
30 switch events. Any method for detecting T cell proliferation known in the art can be used with the immunodominant vespid antigen 5 peptides obtained according to the present invention.

- Thus, in a preferred aspect, the peptides of the present invention can be used in *in vitro* assays with peripheral blood lymphocytes or cell lines derived from peripheral blood lymphocytes, obtained from vespid venom antigen 5 sensitive individuals to detect secretion of lymphokines ordinarily associated with allergic responses, *e.g.*, IL-4. Such assays may indicate which epitopes are responsible or associated with the allergic condition. In this way, specific epitopes responsible for T cell responses associated with allergic response can be identified. The sequences of such epitopes can be compared to other vespid venom antigen 5s and to environmental or autologous proteins to determine if there are sequence similarities that suggest possible cross-reactivity. The peptides can be tested for the ability to induce T cell anergy, *e.g.*, by mega-dose administration, modification to produce an epitope antagonist, administration in the absence of the appropriate costimulatory signals, and other methods thought to result in T cell anergy. Peptides containing such epitopes are ideal candidates for therapeutics.
- In a further embodiment, the polypeptides of the invention can be used directly in assays to detect the extent of cross-reactivity with other environmental proteins and/or homologous proteins, with which they share sequence similarity. In particular, the immunodominant fragments of the vespid venom antigen 5s that have sequence similarity with such environmental, and more particularly, homologous proteins can be evaluated for cross reactivity with antibodies or T cell specific for such proteins. In a specific embodiment, the cross reactivity of vespid venom antigen 5s with human and mouse testes specific protein can be evaluated.

Diagnostic and Therapeutic Uses of the Peptides

- The present invention identifies therapeutically relevant polypeptide fragment of vespid venom antigen 5. The invention contemplates use of the immunoreactive fragments of vespid venom antigen 5 for the preparation of diagnostic or therapeutic compositions, for the use in the diagnosis and therapy of vespid venom allergen-specific allergic conditions. In particular, vespid antigen 5 from *Dolichovespula maculata* (white-face hornet) (*Dol m V*), *Dolichovespula arenaria* (yellow hornet) (*Dol a V*), *Vespula vulgaris* (yellowjacket) (*Ves v V*), *Vespula maculifrons* (yellowjacket) (*Ves m V*), *Polistes annularis* (wasp) (*Pol a V*), and *Polistes exclamans* (wasp) (*Pol e V*) are contemplated for use in diagnosis and therapy according to the present invention. Other vespid species known to harbor antigen 5, and thus represent additional antigen 5 species of the invention, include but are not

limited to *Vespa crabo* (European hornet), *V. flavopilosa* (yellow jacket), *V. germanica* (yellowjacket), *V. pennsylvanica* (yellowjacket), *V. squamosa* (yellowjacket), *V. vidua* (yellowjacket), and *P. fuscatus* (paperwasp).

5

Diagnostic Methods

The present invention contemplates *in vitro* diagnostic assays on peripheral blood lymphocytes, as described *supra*. Such diagnostic assays can give detailed information about the antigen 5-specific T cell responses, the phenotype of the T cell response, and preferably the T cell epitope of the antigen 5 involved in T cell responses. The immunodominant epitope and the epitope involved in IgE isotype class switch events can be detected, if they are not the same. In particular, the T cell epitopes of vespid venom antigen 5s that stimulate proliferation and/or lymphokine secretion of T cells of a phenotype associated with IgE isotype class switching events can be identified for a specific individual, or for a class of individuals who share MHC haplotype or a predominant T cell receptor variable region expression, or both.

In vivo assays for allergenicity generally consist of skin prick sensitivity assays, in which serially diluted amounts of an allergen are administered either subcutaneously or intradermally into a patient's skin, and wheel and erythema reactions are detected. As with *in vitro* assays, the availability of pure venom antigen 5 peptides greatly increases the value of the results of the *in vivo* diagnostic assays since cross-reactivity with impurities in extracts prepared from vespid venom sacs, and uncertainty about the epitope specificity can be avoided.

25

Therapeutic Methods

Therapeutic compositions of the invention (see, *infra*) can be used in immunotherapy, also referred to as hyposensitization therapy. Immunotherapy has proven effective in allergic diseases, particular insect allergy. Allergens are administered parenterally over a long period of time in gradually increasing doses. Such therapy may be particularly effective when the allergen or allergens to which the patient is sensitive have been specifically identified and the therapy is targeted to those allergen(s). Thus, the availability of specific, pure vespid venom antigen 5 peptide(s) in large quantities is important for immunotherapy of allergy.

The present invention contemplates use of peptides containing at least an immunomodulatory T cell epitope of a vespid venom antigen 5 to induce specific T cell anergy to the vespid venom antigen 5. Such immunomodulatory peptide contains one or more immunodominant epitope of antigen 5. A peptide comprising such a T cell epitope and lacking a B cell epitope can be administered to a patient. As discussed *supra*, the presence of B cell epitopes on an allergen can cause an undesirable systemic reaction when the allergen is used for immunotherapy. Thus, a particular advantage of the invention is the capability to provide allergen polypeptides that do not cause undesirable systemic effects.

10

In one embodiment, one or more peptides can be injected subcutaneously to decrease the T cell response to the entire molecule, *e.g.*, as described by Briner et al. [*Proc. Natl. Acad. Sci. U.S.A.*, **90**:7608-12 (1993)]. In another embodiment, one or more peptide can be administered intranasally to suppress allergen-specific responses in naive and sensitized subjects [see *e.g.*, Hoyne et al., *J. Exp. Med.*, **178**:1783-88 (1993)].

15

Administration of a vespid venom antigen peptide of the invention is expected to induce anergy, resulting in cessation of allergen-specific antibody production or allergen-specific T cell response, or both, and thus, have a therapeutic effect.

20

In a preferred aspect of the invention, peptide-based therapy to induce T cell anergy is customized for each individual or a group of individuals. Using the diagnostic methods of the present invention, the specific T cell epitope or epitopes of a vespid venom antigen 5 involved in the allergic response can be identified. Peptides comprising these epitopes can then be used in an individualized immunotherapy regimen.

25

For example, by analogy and based on the data shown in Table 4, *infra*, a treatment protocol for BALB/c mice could include peptides 1, 5, 6, 11, 15, 20, and 18, while treatment of ASW/Sn mice might include peptides 6, 7, and 18; treatment of C3H/He might include peptides 2, 5, and 15; and treatment of P/J mice might include peptides 1, 9, and 15.

30

Administration of the therapeutic compositions of the present invention to desensitize an individual can be carried out using known techniques. For example, one or more

immunodominant peptides of the invention comprising at least one T cell epitope can be administered in combination with an appropriate diluent, a carrier, and/or an adjuvant. To induce T cell anergy in an individual, the therapeutic composition is preferably administered in non-immunogenic form, *e.g.*, it does not contain adjuvant.

- 5 Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Pharmaceutically acceptable carriers include polyethylene glycol [Wie et al., *International Archives of Allergy and Applied Immunology*, **64**:84-99 (1981)] and liposomes [Strejan et al., *Journal of Neuroimmunology*, **7**:27 (1984)]. Such compositions will generally be administered by injection (subcutaneous, intravenous, etc.), oral administration (*e.g.*, as in
- 10 the form of a capsule), transdermal application, or transmucosal administration (*e.g.* nasal, buccal, or rectal administration). The therapeutic compositions of the invention are administered to individuals sensitive to vespid venom at dosages and for lengths of time effective to reduce sensitivity (*i.e.*, reduce the allergic response) of the individual to vespid stings. A therapeutically effective amount of one or more of the same or different
- 15 therapeutic compositions can be administered simultaneously or sequentially to an individual sensitive to vespid venom. Effective amounts of the therapeutic compositions will vary according to factors such as the degree of sensitivity of the individual to vespid venom, the age, sex, and weight of the individual, and the ability of the peptide to stimulate a T cell response in the individual.

20

- In yet another aspect of the present invention, a composition is provided comprising at least two immunodominant peptides (*e.g.*, a physical mixture of at least two peptides), each comprising at least one T cell epitope of antigen 5. The peptides are derived from the same or from different homologues of antigen 5. Such compositions can be
- 25 administered in the form a therapeutic composition with a pharmaceutically acceptable carrier or diluent. A therapeutically effective amount of one or more of such compositions can be administered simultaneously or sequentially to an individual sensitive to vespid venom.

- 30 **Transmucosal administration.** According to the invention, any transmucosal route of administration, including but not limited to nasal, rectal, oral, vaginal, buccal, etc. can be employed. It can be readily appreciated that any of the transmucosal routes of administration may be enhanced by use of a mucosal penetration enhancer. The term "mucosal penetration enhancer" refers to a reagent that increases the rate or facility of

transmucosal penetration of peptide, such as but not limited to, a bile salt, fatty acid, surfactant or alcohol. In specific embodiments, the permeation enhancer can be sodium cholate, sodium dodecyl sulphate, sodium deoxycholate, taurodeoxycholate, sodium glycocholate, dimethylsulfoxide or ethanol. Suitable penetration enhancers also include
5 glycyrrhetic acid [U.S. Patent No. 5,112,804 to Kowarski] and polysorbate-80, the latter preferably in combination with a non-ionic surfactant such as nonoxynol-9, laureth-9, poloxamer-124, octoxynol-9, or lauramide-DEA [European Patent EP 0 242 643 B1 by Stoltz]. The selection of a particular mucosal penetration enhancer may depend on the characteristics of the specific mucosa. These factors are addressed in greater detail below.

10

Nasal Administration. Transnasal or intranasal (nasal) delivery of an antigen 5 peptide can be used to suppress an allergen-specific response [see Hoyne et al., *J. Exp. Med.*, **178**:1783-1788 (1993)]. Nasal delivery allows the passage of the protein to the blood stream directly after administering the product to the nose, without the necessity for
15 deposition of the product in the lung. Formulations for nasal delivery include those with dextran or cyclodextran. For nasal administration, a useful device is a small, hard bottle to which a metered dose sprayer is attached. In one embodiment, the metered dose is delivered by drawing the peptide solution into a chamber of defined volume, which chamber has an aperture dimensioned to aerosolize and aerosol formulation by forming a
20 spray when a liquid in the chamber is compressed. The chamber is compressed to administer the solution. In a specific embodiment, the chamber is a piston arrangement. Such devices are commercially available. Alternatively, a plastic squeeze bottle with an aperture or opening dimensioned to aerosolize an aerosol formulation by forming a spray when squeezed. The opening is usually found in the top of the bottle, and the top is
25 generally tapered to partially fit in the nasal passages for efficient administration of the aerosol formulation. Preferably, the nasal applicator will provide a metered amount of the nasal formulation, for administration of a measured dose of the peptide(s).

Administration Via Suppositories. In another aspect, peptide is formulated in a matrix
30 suitable for rectal (or vaginal) insertion, *i.e.*, in a suppository. The invention is not limited to any particular suppository formulation. Indeed, many suppository formulations are known in the art, *e.g.*, as described in *Remington's Pharmaceutical Sciences*, *Physician's Desk Reference*, and *U.S. Pharmacopeia*.

Administration Via a Buccal Patch. According to the invention, peptide can be formulated in a buccal patch for administration via the interior of the cheek. It may be appreciated that a buccal patch constitutes another form of transmucosal administration. The technology for preparing buccal patch formulations is known in the art, *e.g.*, *Remington's*
5 *Pharmaceutical Sciences, supra*.

Oral-Pharyngeal Administration. In yet another embodiment, peptide can be formulated for oral-pharyngeal, including sublingual and transbuccal, administration. For example, peptide can be incorporated in a "candy" matrix, such as that described in U.S. Patent
10 No. 4,671,953, in a gum base, or a lozenge. In another embodiment, the peptide can be formulated in a capsule or pill form for sublingual placement. It is particularly contemplated that peptide for oral-pharyngeal administration may be formulated with a flavor masking agent or coating. Many flavor masking agents for use with oral
15 pharmaceuticals are known in the art, and can be selected for use with the present invention.

Oral Administration. In still a further embodiment, peptide can be formulated for oral administration via the stomach and intestinal mucosa. For oral administration, peptide can be administered in a carrier designed for drug release in either the stomach (an acidic
20 environment), or the intestines, or both. Many capsules, pills, and matrices for oral administration of a drug are known in the art, and can be selected on the basis of compatibility with peptide, and the desired point and rate of drug release by the ordinary skilled physician. Oral administration of peptide may require higher dosages than other routes of administration to overcome the effects of first pass metabolism by the liver.

25
Transdermal Administration. In a further embodiment, as noted above, the present invention is directed to transdermal administration of peptide. Various and numerous methods are known in the art for transdermal administration of a compound, *e.g.*, via a transdermal patch. These methods and associated devices provide for control of the rate
30 and quantity of administration of a drug, and some allow for continuous modulation of drug delivery. Transdermal patches are described in, for example, U.S. Patent No. 5,407,713, issued April 18, 1995 to Rolando et al.; U.S. Patent No. 5,352,456, issued October 4, 1994 to Fallon et al.; U.S. Patent No. 5,332,213 issued August 9, 1994 to D'Angelo et al.; U.S. Patent No. 5,336,168, issued August 9, 1994 to Sibalis; U.S.

Patent No. 5,290,561, issued March 1, 1994 to Farhadieh et al.; U.S. Patent No. 5,254,346, issued October 19, 1993 to Tucker et al.; U.S. Patent No. 5,164,189, issued November 17, 1992 to Berger et al.; U.S. Patent No. 5,163,899, issued November 17, 1992 to Sibalis; U.S. Patent Nos. 5,088,977 and 5,087,240, both issued February 18, 5 1992 to Sibalis; U.S. Patent No. 5,008,110, issued April 16, 1991 to Benecke et al.; and U.S. Patent No. 4,921,475, issued May 1, 1990 to Sibalis, the disclosure of each of which is incorporated herein by reference in its entirety.

It can be readily appreciated that a transdermal route of administration may be enhanced 10 by use of a dermal penetration enhancer, *e.g.*, such as enhancers described in U.S. Patent No. 5,164,189 (*supra*), U.S. Patent No. 5,008,110 (*supra*), and U.S. Patent No. 4,879,119, issued November 7, 1989 to Aruga et al., the disclosure of each of which is incorporated herein by reference in its entirety.

15 Pharmaceutically Acceptable Compositions

The *in vivo* diagnostic or therapeutic compositions of the invention may also contain appropriate pharmaceutically acceptable carriers, excipients, diluents and adjuvants. As used herein, the term "pharmaceutically acceptable" preferably means approved by a regulatory agency of a government, in particular the Federal government or a state 20 government, or listed in the U.S. Pharmacopeia or another generally recognized pharmacopeia for use in animals, and more particularly in humans, although a pharmaceutically acceptable carrier may share the attributes of such approved or recognized carriers without having itself been approved or recognized. Examples of suitable pharmaceutical carriers are provided in "Remington's Pharmaceutical Sciences" 25 by E.W. Martin.

Such pharmaceutically acceptable carriers can be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame oil and the like. Water is a preferred carrier when the 30 pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include mannitol, human serum albumin (HSA), starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, magnesium carbonate, magnesium stearate, sodium stearate, glycerol monostearate,

talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. These compositions can take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained-release formulations and the like.

- 5 Such compositions will contain an effective diagnostic or therapeutic amount of the active compound together with a suitable amount of carrier so as to provide the form for proper administration to the patient. While intravenous injection is a very effective form of administration, other modes can be employed, such as by injection, or by oral, nasal, or parenteral administration.

10

The invention will be further clarified by the following examples, which are intended to be purely exemplary of the invention.

15 EXAMPLE 1: MAPPING T CELL EPITOPES OF DOL M Ag5

- In the present Example, a series of 15 to 20-residue peptides with 10-residue overlaps to encompass the entire white faced hornet Ag5 molecule (form 2) was synthesized. The sequences of these peptides are given in Figure 2. These peptides were used for mapping
- 20 T cell epitopes of hornet Ag5 by their stimulation of specific murine spleen cells immunized with native or recombinant Antigen 5. These peptides were also used to study the cross reacting T cell epitopes of hornet antigen 5 and a mouse testis protein Tpx [Mizuku et al., *Mammalian Genome*, 3:274-280 (1992)]. This cross reactivity is suggested by the partial sequence identity of hornet antigen 5 with mouse and human testis
- 25 proteins (Figure 3).

Results

- Antibody response in mice of different haplotypes to recombinant hornet Ag5.* Five strains of mice were immunized with recombinant (r) white faced hornet Ag5 and alum as
- 30 adjuvant. Four strains were found to be high antibody responders for r-white faced hornet Ag5: BALB/v byj, ASW/Sn, C3H/He, and P/J. Antibody responses were evaluated by direct ELISA [King et al., *J. Immunol.*, 154:577 (1995)]. The antibody titers (the reciprocal of the dilution yielding 50% of maximum signal extrapolated from ELISA curves in wells coated with r-Dol m 5.02 or n-Dol m 5.01) of these sera for n-(natural)

and r-Ag5 are given in Table 3. One strain, C57Bl/6, was found to be a poor antibody responder.

Table 3

5

**Antibody for Mice of Different Strains
Immunized with Recombinant White faced hornet Ag5**

Mice		Half maximal Ab titer*	
Strain	Haplotype	r-Dol m 5.02	n-Dol m 5.01
10 C57Bl/6	b	1×10^2	2×10^1
BALB/c byj	d	8×10^4	4×10^4
C3H/He	k	6×10^3	2×10^3
P/J	p	2×10^4	1×10^4
15 ASW/Sn	s	1×10^5	1×10^5

*Sera are from week five bleeding after three intraperitoneal immunizations with 0.2 ml of 10 μ g/ml r-Dol m 5.02 and alum (5 μ g/ml) in 0.05 M sodium phosphate buffer, pH 6.2, on weeks 0, 2, and 4.

20 *T cell response in BALB/c mice to natural hornet Ag5 and its recombinant protein or fragments.* Figures 4 and 5 show the proliferative responses of spleen cells from BALB/c mice immunized 4 times with n- and r-Ag5 respectively. Nearly identical results show that r-Ag5 and n-Ag5 share common continuous T cell epitopes. Previous studies have shown that r-Ag5 lacks the discontinuous B cell epitopes of n-Ag5, as r-Ag5 is not properly folded with the disulfide bonds of n-Ag5 [King et al., *J. Immunol.*, **154**:577-584 (1995)]. Figure 6 shows the response of spleen cells from BALB/c mice immunized 5 times with r-Ag5. Higher proliferative responses were observed than those in Figure 5, where mice were immunized 4 times with r-Ag5. The results in Figure 4-6 together indicate that peptides 1, 5, 6, 11, 15, 20, and 18 contain the T cell epitopes of Ag5, if we assume that peptides have stimulation indices of ≥ 4 represent positive responses with a high degree of certainty.

Figures 7 and 8 show the proliferation data with spleen cells from mice immunized 4 times with recombinant N-terminal and C-terminal fragments of antigen 5, labelled IN (residue 1-114) and C (residue 151-204), respectively. Similar but higher proliferation results were obtained with mice after 5 times immunization, as given in Figure 9 and 10. As to be expected, spleen cells from r-fragment IN or C immunized mice were stimulated by peptides whose sequences are represented by the immunogen. There is one exception

shown in Figure 9: spleen cells from r-fragment IN (residue 1-114) immunized mice were stimulated by peptide 18 (residue 176-195).

T cell response in mice of different haplotypes to recombinant hornet Ag5. Figures 11 to 13 show the proliferative response of spleen cells for mice of ASW/Sn, P/J, and C3H/He, respectively. The results for ASW/Sn mice in Figure 11 indicate that there are four peptides with stimulation indices of ≥ 4 : peptides 6, 7, 9, and 18. The results for P/J mice in Figure 12 show multiple peptides with stimulation indices of ≥ 4 : peptides 1, 2, 5, 9, 12, 13, 14, 15, 20, and 17. The data for C3H/He mice in Figure 12 indicate three peptides with stimulation indices of ≥ 4 : peptides 2, 5 and 15.

Figure 14 shows the proliferation responses of spleen cells for C57Bl/6 mice. Several peptides had stimulation indices of ≥ 2 , but only one peptide, number 9, showed a stimulation index approaching 4. A lower proliferation response compared to the other mouse strains was also observed with r-Ag5 as a control. These data indicate that C57Bl/6 mice showed poor T cell responses to white faced hornet Ag5, as contrasted to the other 4 strains of mice tested (Table 4).

Table 4

Summary of proliferation indices of r-Ag5 specific spleen cells from mice of different strains on stimulation with the dominant T cell epitope peptides¹

5	Stimulating ² antigen	Stimulation Index ³				
		BALB/c	ASW/Sn	C3H/He	P/J	C57Bl/6
10	Peptide 1	8.6	1.9	2.4	6.9	2.5
	2	2.9	2.2	4.3	4.8	1.9
	5	9.3	1.4	5.8	4.0	1.7
	6	5.8	8.8	2.0	2.2	1.9
	7	(0.9)	7.7	2.2	3.3	2.1
15	9	3.8	4.0	2.7	9.9	3.9
	11	5.8	0.9	3.0	1.4	2.7
	12	2.5	1.4	3.3	4.0	2.5
	13	(1.3)	1.2	3.6	4.5	2.5
	14	2.3	2.3	2.9	4.2	2.0
20	15	5.6	1.4	4.7	7.8	2.8
	20	6.5	1.7	2.8	4.5	2.1
	17	(1.1)	1.6	3.6	4.1	2.4
	18	17.7	9.1	2.9	2.5	2.7
	r-Ag5	20.9	15.4	12.3	20.8	7.0
25	Blank	3970 cpm	5640 cpm	3130 cpm	1730 cpm	1050 cpm

NOTES

1. Dominant T cell epitopes are defined as those peptides which showed stimulation indices of ≥ 4 in at least one of 5 strains of mice tested.
2. Peptides in bold face represent those with stimulation indices of ≥ 4 in 2 or more strains of mice.
3. Stimulation index values are taken from Figure 5, 9, 10, 11 and 12, in which spleen cells were from mice immunized 5 times with r-Ag5. The exceptions are those values in parentheses which are taken from Figure 4 in which spleen cells were from 4 times r-Ag5 immunized mice.

35

T cell epitopes of hornet Ag5 and a mouse testis protein Tpx. Cross reactivity of these two proteins was studied in mice immunized with their fragments (Figures 15 and 16). Spleen cells from BALB/c mice immunized with r-fragment Tpx-N (residue 14-106) responded equally well to stimulation with the immunogen or the r-fragment IN of hornet Ag5 (residue 1-114). Similarly spleen cells from BALB/c mice immunized with r-fragment Tpx-C (residue 101-163) responded equally well to stimulation with the immunogen or the r-fragment C of hornet Ag5 (residue 151-204).

40

Proliferation of Tpx-N or -C primed spleen cells was also tested on stimulation with synthetic peptides which correspond to sequences present in r-fragment IN and C of hornet Ag5. The data in Figure 15 show high stimulation indices (greater than 6) for both r-fragments Tpx-N and fragment IN, only moderate stimulation indices (about 3) for peptides 7, 8 and 9, and near baseline stimulation indices of about 2 or less for the remaining peptides. The data in Figure 16 show high stimulation indices (greater than 15) for Tpx-C, 9 for fragment C, 6 for peptide 18, and 2 or less for all other peptides.

The data in Figure 15 and 16 were obtained from mice after 4 immunizations. Further studies after five immunizations are planned as the data in Figures 7 to 10 indicate that stronger stimulation indices are obtained after longer immunization.

Discussion

The present Example defines indices of 4 or greater for proliferation of Ag5- specific cells as containing a T cell epitope of Ag5. With this definition, the data in Figures 3-13 for mice of 5 strains with different haplotypes would indicate that the T cell epitopes of Ag5 are distributed throughout the entire molecule. This is summarized in Table 4. As C57Bl/6 mice is a poor responder, its data are excluded in the following comparison.

The data in Table 4 indicate that 14 peptides contain T cell epitopes of Ag5. Three of these peptides are recognized by three of the four strains tested, five of them by two strains and the remaining six by only one strain. The peptides which are recognized by two or three strains of mice are indicated in bold face characters in Table 2. They are peptides 1, 2, 5, 6, 9, 15, 20 and 18. As can be seen in Figure 1, there are extensive sequence identities of these white faced hornet peptides with those of the other vespids. This is particularly apparent for peptides 6, 20, and 18.

In addition to the seven vespid Ag5 sequences in Figure 1, there are eight other vespid Ag5s with known sequences. They are nearly identical to those in Figure 1, depending on their species group [Lu, et al., *J. Immunol.*, **150**:2823-2830 (1993); Hoffman, *J. Allergy Clin. Immunol.*, **92**:707-716 (1993)]. Fire ant venom allergen Sol i 3 also has sequence similarity with vespid Ag5s [Hoffman, *J. Allergy Clin. Immunol.*, **91**:71-78 (1993)]. Peptides 9, 20 and 18 of white faced hornet Ag5 have a high degree of sequence identity with the corresponding peptides of Sol i 3.

It is most likely that these vespid and fire ant Ag5s have T cell epitopes of identical and/or similar sequences. These epitopes studies are being continued with yellowjacket or paper wasp Ag5 immunized mice.

- 5 The finding of cross reactive T cell epitopes of hornet Ag5 and a mammalian testis protein is interesting, although its significance in clinical allergy, if any, is unknown. Published reports do not indicate any unusual distribution of male and female insect allergic patients. Our own unpublished studies showed that male and female BALB/c mice gave indistinguishable antibody response when immunized with hornet Ag5 in presence of
10 alum.

- The present invention is not to be limited in scope by the specific embodiments describe herein. Indeed, various modifications of the invention in addition to those described
15 herein will become apparent to those skilled in the art from the foregoing description and the accompanying figures. Such modifications are intended to fall within the scope of the appended claims.

- It is further to be understood that all base sizes or amino acid sizes, and all molecular
20 weight or molecular mass values, given for nucleic acids or polypeptides are approximate, and are provided for description.

Various publications are cited herein, the disclosures of which are incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

- 1 1. A peptide characterized by the following properties:
- 2 a) having between 8 and 35 amino acid residues of vespid venom antigen 5;
- 3 and
- 4 b) it is antigenic for T cell proliferation in a mouse immunized with a vespid
- 5 venom antigen 5, which mouse is a strain selected from the group consisting or
- 6 BALB/c, ASW/Sn, C3H/He, and P/J.

- 1 2. The peptide of claim 1 which corresponds to a fragment of white face hornet
- 2 antigen 5, form 2, selected from the group consisting of:

3	NNYCKIKCRKGIHTLCKFGT	(SEQ ID NO:8);
4	GIHTLCKFGTSMKPNCGRNV	(SEQ ID NO:9);
5	KNEILKRHNDFRQNVAKGLE	(SEQ ID NO:12);
6	FRQNVAKGLETRGKPGPQPP	(SEQ ID NO:13);
7	TRGKPGPQPPAKNMNVLVWN	(SEQ ID NO:14);
8	DELA KIAQTWANQCDFNHDD	(SEQ ID NO:16);
9	CRNTAKYQVGQNIAISSTA	(SEQ ID NO:18);
10	QNIAISSTTATQFDRPSKLI	(SEQ ID NO:19);
11	TQFDRPSKLIKQWEDEVTEF	(SEQ ID NO:20);
12	KQWEDEVTEFNYKVGLQNSN	(SEQ ID NO:21);
13	NYKVGLQNSNFRKVGHYTQM	(SEQ ID NO:22);
14	FRKVGHYTQMVWGKT	(SEQ ID NO:23);
15	KEIGCGSIKYIEDN WYTHYL	(SEQ ID NO:25); and
16	IEDN WYTHYLVCNYGPGGND	(SEQ ID NO:26).

- 1 3. The peptide of claim 1 which corresponds to a fragment of white face hornet
- 2 antigen 5, form 2, selected from the group consisting of:

3	NNYCKIKCRKGIHTLCKFGT	(SEQ ID NO:8);
4	GIHTLCKFGTSMKPNCGRNV	(SEQ ID NO:9);
5	KNEILKRHNDFRQNVAKGLE	(SEQ ID NO:12);
6	FRQNVAKGLETRGKPGPQPP	(SEQ ID NO:13);
7	DELA KIAQTWANQCDFNHDD	(SEQ ID NO:16);
8	NYKVGLQNSNFRKVGHYTQM	(SEQ ID NO:22);

9 FRKVGHYTQMVGKT (SEQ ID NO:23); and
 10 IEDNWYTHYLCNYGPGGND (SEQ ID NO:26).

1 4. The peptide of claim 1 having the amino acid sequence
 2 X_1 BYCKIX₂CX₃X₄GX₅X₆HTX₇CX₈X₉G (SEQ ID NO:31), wherein X₁ is a neutral amino
 3 acid residue, X₂ is a basic amino acid residue or deleted, X₃ is any amino acid residue, X₄
 4 is a polar amino acid, X₅ is glycine or deleted, X₆ is any amino acid residue, X₇ is an
 5 amino acid residue with an aliphatic side chain, X₈ is a polar amino acid residue, and X₉
 6 is an amino acid residue with an aromatic side chain.

1 5. The peptide of claim 4, wherein X₁ is valine or asparagine; X₂ is lysine, arginine,
 2 or deleted; X₃ is proline, arginine, serine, or leucine; X₄ is arginine, lysine, or serine; X₆
 3 isoleucine, leucine, threonine, or valine; X₇ is alanine, valine, isoleucine, or leucine; X₈ is
 4 lysine, arginine, glutamine, or asparagine; or X₉ is phenylalanine, tryptophan, or tyrosine;
 5 or any combination of the foregoing.

1 6. The peptide of claim 1 having the amino acid sequence
 2 GX₅X₆HTX₇CX₈X₉GX₁₀SX₁₁KPX₁₂X₁₃NCX₁₄X₁₅X₁₆X₁₇X₁₈ (SEQ ID NO:32), wherein X₅ is
 3 glycine or deleted, X₆ is any amino acid residue, X₇ is an amino acid residue with an
 4 aliphatic side chain, X₈ is a polar amino acid residue, X₉ is an amino acid residue with an
 5 aromatic side chain, X₁₀ is a polar amino acid residue, X₁₁ is any amino acid residue, X₁₂
 6 is a polar amino acid residue or deleted, X₁₃ is a basic amino acid residue or deleted, X₁₄
 7 is a small chain amino acid residue, X₁₅ is any amino acid residue, X₁₆ is a basic or polar
 8 neutral amino acid residue, X₁₇ is any amino acid residue, and X₁₈ is an aliphatic amino
 9 acid residue.

1 7. The peptide of claim 6, wherein X₆ isoleucine, leucine, threonine, or valine; X₇ is
 2 alanine, valine, isoleucine, or leucine; X₈ is lysine, arginine, glutamine, or asparagine; X₉
 3 is phenylalanine, tryptophan, or tyrosine; X₁₀ is threonine, serine, aspartic acid, or
 4 glutamic acid; X₁₁ is leucine, isoleucine, methionine, or threonine; X₁₂ is serine or
 5 deleted; X₁₃ is lysine, arginine, or deleted; X₁₄ is glycine or alanine; X₁₅ is asparagine,
 6 glutamine, glycine, serine, arginine, or lysine; X₁₆ is lysine, arginine, asparagine, or
 7 glutamine; X₁₇ is lysine, arginine, isoleucine, leucine, or valine; or X₁₈ is alanine, valine,
 8 isoleucine, or leucine; or any combination of the foregoing.

1 8. The peptide of claim 1 having the amino acid sequence
 2 X_1 BYCKIX₂CX₃X₄GX₅X₆HTX₇CX₈X₉GX₁₀SX₁₁KPX₁₂X₁₃NCX₁₄X₁₅X₁₆X₁₇X₁₈ (SEQ ID
 3 NO:33), wherein X_1 is a neutral amino acid residue, X_2 is a basic amino acid residue or
 4 deleted, X_3 is any amino acid residue, X_4 is a polar amino acid, X_5 is glycine or deleted,
 5 X_6 is any amino acid residue, X_7 is an amino acid residue with an aliphatic side chain, X_8
 6 is a polar amino acid residue, X_9 is an amino acid residue with an aromatic side chain, X_{10}
 7 is a polar amino acid residue, X_{11} is any amino acid residue, X_{12} is a polar amino acid
 8 residue or deleted, X_{13} is a basic amino acid residue or deleted, X_{14} is a small chain amino
 9 acid residue, X_{15} is any amino acid residue, X_{16} is a basic or polar neutral amino acid
 10 residue, X_{17} is any amino acid residue, and X_{18} is an aliphatic amino acid residue.

1 9. The peptide of claim 8, wherein X_1 is valine or asparagine; X_2 is lysine, arginine,
 2 or deleted; X_3 is proline, arginine, serine, or leucine; X_4 is arginine, lysine, or serine; X_6
 3 isoleucine, leucine, threonine, or valine; X_7 is alanine, valine, isoleucine, or leucine; X_8 is
 4 lysine, arginine, glutamine, or asparagine; X_9 is phenylalanine, tryptophan, or tyrosine;
 5 X_{10} is threonine, serine, aspartic acid, or glutamic acid; X_{11} is leucine, isoleucine,
 6 methionine, or threonine; X_{12} is serine or deleted; X_{13} is lysine, arginine, or deleted; X_{14}
 7 is glycine or alanine; X_{15} is asparagine, glutamine, glycine, serine, arginine, or lysine; X_{16}
 8 is lysine, arginine, asparagine, or glutamine; X_{17} is lysine, arginine, isoleucine, leucine, or
 9 valine; or X_{18} is alanine, valine, isoleucine, or leucine; or any combination of the
 10 foregoing.

1 10. The peptide of claim 1 having the amino acid sequence
 2 $KX_{19}X_{20}IX_{21}X_{22}X_{23}HNX_{24}FRQKX_{25}AX_{26}GLE$ (SEQ ID NO:34), wherein X_{19} is a basic or
 3 neutral polar amino acid residue X_{20} is any amino acid residue X_{21} is an aliphatic amino
 4 acid residue, X_{22} is a polar amino acid residue, X_{23} is a polar charged amino acid residue,
 5 X_{24} is a polar amino acid residue, X_{25} is an aliphatic amino acid residue; and X_{26} is a polar
 6 basic or neutral amino acid residue.

1 11. The peptide of claim 10, wherein X_{19} is asparagine, glutamine, or lysine; X_{20} is
 2 aspartic acid, glutamic acids, leucine, or isoleucine; X_{21} is valine, leucine, or isoleucine;
 3 X_{22} arginine, asparagine, or serine; X_{23} is glutamic acid or arginine; X_{24} is aspartic acid,
 4 glutamic acid, asparagine, glutamine, or arginine; X_{25} is valine, leucine, or isoleucine; or
 5 X_{26} is arginine, asparagine, or lysine; or any combination of the foregoing.

- 1 12. The peptide of claim 1 having the amino acid sequence
2 FRQKX₂₅AX₂₆GLETRGX₂₇PGPQPX₂₈ (SEQ ID NO:35), wherein X₂₅ is an aliphatic amino
3 acid residue, X₂₆ is a polar basic or neutral amino acid residue, X₂₇ is a polar basic or
4 neutral amino acid residue, and X₂₈ is any amino acid residue.
- 1 13. The peptide of claim 12, wherein X₂₅ is valine, leucine, or isoleucine; X₂₆ is
2 arginine, asparagine, or lysine; X₂₇ is asparagine or lysine; or X₂₈ is glycine, alanine, or
3 proline; or any combination of the foregoing.
- 1 14. The peptide of claim 1 having the amino acid sequence
2 KX₁₉X₂₀IX₂₁X₂₂X₂₃HNX₂₄FRQKX₂₅AX₂₆GLETRGX₂₇PGPQPX₂₈ (SEQ ID NO:36), wherein
3 X₁₉ is a basic or neutral polar amino acid residue X₂₀ is any amino acid residue X₂₁ is an
4 aliphatic amino acid residue, X₂₂ is a polar amino acid residue, X₂₃ is a polar charged
5 amino acid residue, X₂₄ is a polar amino acid residue, X₂₅ is an aliphatic amino acid
6 residue, X₂₆ is a polar basic or neutral amino acid residue, X₂₇ is a polar basic or neutral
7 amino acid residue, and X₂₈ is any amino acid residue.
- 1 15. The peptide of claim 14, wherein X₁₉ is asparagine, glutamine, or lysine; X₂₀ is
2 aspartic acid, glutamic acids, leucine, or isoleucine; X₂₁ is valine, leucine, or isoleucine;
3 X₂₂ arginine, asparagine, or serine; X₂₃ is glutamic acid or arginine; X₂₄ is aspartic acid,
4 glutamic acid, asparagine, glutamine, or arginine; X₂₅ is valine, leucine, or isoleucine; X₂₆
5 is arginine, asparagine, or lysine; X₂₇ is asparagine or lysine; or X₂₈ is glycine, alanine, or
6 proline; or any combination of the foregoing.
- 1 16. The peptide of claim 1 having the amino acid sequence
2 DELAX₂₉X₃₀AQX₃₁WAX₃₂QCX₃₃X₃₄X₃₅X₃₆HD (SEQ ID NO:37), wherein X₂₉ is any
3 amino acid residue, X₃₀ is an aliphatic amino acid residue, X₃₁ is any amino acid residue,
4 X₃₂ is a polar neutral amino acid residue, X₃₃ is a polar neutral or acidic amino acid
5 residue, X₃₄ is an aromatic amino acid residue, X₃₅ is an aliphatic amino acid residue or
6 deleted, and X₃₆ is any amino acid residue.
- 1 17. The peptide of claim 16, wherein X₂₉ is tyrosine, lysine, or histidine; X₃₀ is
2 isoleucine, leucine, or valine; X₃₁ is threonine or valine; X₃₂ is serine or asparagine; X₃₃ is
3 glutamine, asparagine, aspartic acid, or serine; X₃₄ is phenylalanine to tyrosine; X₃₅ is

- 4 isoleucine, leucine, or deleted; or X_{36} is glycine, asparagine, or valine; or any
5 combination of the foregoing.

1 18. The peptide of claim 1 having the amino acid sequence

2 $NX_{37}X_{38}X_{39}X_{40}X_{41}X_{42}X_{43}X_{44}BX_{45}FX_{46}KX_{47}GHYTOTM$ (SEQ ID NO:38), wherein X_{37} is a
3 cyclic amino acid residue, X_{38} is a polar basic or neutral amino acid residue, X_{39} is any
4 amino acid residue, X_{40} is any amino acid residue, X_{41} is a non-polar amino acid residue,
5 X_{42} is any amino acid residue, X_{43} is a polar amino acid residue or glycine, X_{44} is a polar
6 basic or neutral amino acid residue, X_{45} is a polar neutral amino acid or deleted, X_{46} is
7 any amino acid residue, and X_{47} is a non-polar amino acid residue.

1 19. The peptide of claim 18, wherein X_{37} is proline or tyrosine; X_{38} is asparagine,
2 arginine, or histidine; X_{39} is lysine, threonine, or valine; X_{40} is aspartic acid, glycine, or
3 lysine; X_{41} is isoleucine, leucine, phenylalanine, or threonine; X_{42} is glutamine, isoleucine,
4 leucine, methionine, serine, or threonine; X_{43} is asparagine, glutamic acid, histidine,
5 lysine, or glycine, X_{44} is aspartic acid, asparagine, glutamine, or serine; X_{45} is asparagine
6 or deleted, X_{46} is alanine, arginine, leucine, isoleucine, or serine; or X_{47} is isoleucine,
7 leucine, threonine, or valine; or any combination of the foregoing.

1 20. The peptide of claim 1 having the amino acid sequence

2 $FX_{46}KX_{47}GHYTOTMVWX_{48}X_{49}T$ (SEQ ID NO:39), wherein X_{46} is any amino acid residue,
3 X_{47} is a non-polar amino acid residue, X_{48} is a small side chain amino acid residue, and
4 X_{49} is a polar basic or neutral amino acid residue.

1 21. The peptide of claim 20, wherein X_{46} is alanine, arginine, leucine, isoleucine, or
2 serine; X_{47} is isoleucine, leucine, threonine, or valine; X_{48} is glycine or alanine; or X_{49} is
3 lysine or asparagine; or any combination of the foregoing.

1 22. The peptide of claim 1 having the amino acid sequence

2 $NX_{37}X_{38}X_{39}X_{40}X_{41}X_{42}X_{43}X_{44}BX_{45}FX_{46}KX_{47}GHYTOTMVWX_{48}X_{49}T$ (SEQ ID NO:40),
3 wherein X_{37} is a cyclic amino acid residue, X_{38} is a polar basic or neutral amino acid
4 residue, X_{39} is any amino acid residue, X_{40} is any amino acid residue, X_{41} is a non-polar
5 amino acid residue, X_{42} is any amino acid residue, X_{43} is a polar amino acid residue or
6 glycine, X_{44} is a polar basic or neutral amino acid residue, X_{45} is a polar neutral amino

7 acid or deleted, X_{46} is any amino acid residue, X_{47} is a non-polar amino acid residue, X_{48}
 8 is a small side chain amino acid residue, and X_{49} is a polar basic or neutral amino acid
 9 residue.

1 23. The peptide of claim 22, wherein X_{37} is proline or tyrosine; X_{38} is asparagine,
 2 arginine, or histidine; X_{39} is lysine, threonine, or valine; X_{40} is aspartic acid, glycine, or
 3 lysine; X_{41} is isoleucine, leucine, phenylalanine, or threonine; X_{42} is glutamine, isoleucine,
 4 leucine, methionine, serine, or threonine; X_{43} is asparagine, glutamic acid, histidine,
 5 lysine, or glycine, X_{44} is aspartic acid, asparagine, glutamine, or serine; X_{45} is asparagine
 6 or deleted, X_{46} is alanine, arginine, leucine, isoleucine, or serine; X_{47} is isoleucine,
 7 leucine, threonine, or valine; X_{48} is glycine or alanine; or X_{49} is lysine or asparagine; or
 8 any combination of the foregoing.

1 24. The peptide of claim 1 having the amino acid sequence
 2 $X_{50}ZX_{51}X_{52}X_{53}X_{54}X_{55}HYLX_{56}CNYGPX_{57}GNX_{58}X_{59}X_{60}$ (SEQ ID NO:40), wherein X_{50} is a
 3 non-polar amino acid, X_{51} is a polar acidic or neutral amino acid residue, X_{52} is a polar
 4 basic or neutral amino acid residue, X_{53} is a non-polar amino acid residue, X_{54} is a
 5 moderately polar amino acid residue, X_{55} is a polar basic or neutral amino acid residue,
 6 X_{56} is an aliphatic amino acid residue, X_{57} is any amino acid residue, X_{58} is any amino
 7 acid residue, X_{59} is any amino acid residue, and X_{60} is any amino acid residue.

1 25. The peptide of claim 24, wherein X_{50} is isoleucine, leucine, or methionine; X_{51} is
 2 asparagine, aspartic acid, glutamine, or glutamic acid; X_{52} is asparagine or lysine; X_{53} is
 3 methionine or tryptophan; X_{54} glutamine, histidine, or tyrosine; X_{55} is asparagine, lysine,
 4 or threonine; X_{56} isoleucine, leucine, or valine; X_{57} is alanine, glycine, or serine; X_{58} is
 5 aspartic acid, phenylalanine, or tyrosine; X_{59} is glutamine, leucine, isoleucine, methionine,
 6 or phenylalanine; or X_{60} is asparagine, aspartic acid, or glycine; or any combination of the
 7 foregoing.

1 26. A peptide having an amino acid sequence selected from the group consisting of:
 2 a) NNYCKIKCRKGIHTLCKFGT (SEQ ID NO: 8), or its homologue;
 3 b) GIHTLCKFGTSMKPNCGRNV (SEQ ID NO:9), or its homologue;
 4 c) NNYCKIKCRKGIHTLCKFGTGTSMKPNCGRNV (SEQ ID NO:42), or
 5 its homologue;

- 6 d) KNEILKRHND~~F~~RQNVAKGLE (SEQ ID NO:12), or its homologue;
- 7 e) FRQNVAKGLETRGKPGPQPP (SEQ ID NO:13), or its homologue;
- 8 f) KNEILKRHND~~F~~RQNVAKGLETRGKPGPQPP (SEQ ID NO:43), or its
- 9 homologue;
- 10 g) DELAKIAQTTWANQCDFNHDD (SEQ ID NO:16), or its homologue;
- 11 h) NYKVGLQNSNFRKVGHYTQM (SEQ ID NO:22), or its homologue;
- 12 i) FRKVGHYTQMVWGKT (SEQ ID NO:23), or its homologue;
- 13 j) NYKVGLQNSNFRKVGHYTQMVWGKT (SEQ ID NO:44), or its
- 14 homologue; and
- 15 k) IEDNWYTHYLVCNYGPGGND (SEQ ID NO:26), or its homologue.

1 27. A recombinant polypeptide comprising two or more peptides non-contiguously
 2 arranged relative to the native sequence of vespid venom antigen 5, wherein the peptides
 3 are selected from the group consisting of:

- 4 a) NNYCKIKCRKGIHTLCKFGT (SEQ ID NO: 8), or its homologue;
- 5 b) GIHTLCKFGTSMKPNCGRNV (SEQ ID NO:9), or its homologue;
- 6 c) NNYCKIKCRKGIHTLCKFGTGTSMPKNCGRNV (SEQ ID NO:42), or
- 7 its homologue;
- 8 d) KNEILKRHND~~F~~RQNVAKGLE (SEQ ID NO:12), or its homologue;
- 9 e) FRQNVAKGLETRGKPGPQPP (SEQ ID NO:13), or its homologue;
- 10 f) KNEILKRHND~~F~~RQNVAKGLETRGKPGPQPP (SEQ ID NO:43), or its
- 11 homologue;
- 12 g) DELAKIAQTTWANQCDFNHDD (SEQ ID NO:16), or its homologue;
- 13 h) NYKVGLQNSNFRKVGHYTQM (SEQ ID NO:22), or its homologue;
- 14 i) FRKVGHYTQMVWGKT (SEQ ID NO:23), or its homologue;
- 15 j) NYKVGLQNSNFRKVGHYTQMVWGKT (SEQ ID NO:44), or its
- 16 homologue; and
- 17 k) IEDNWYTHYLVCNYGPGGND (SEQ ID NO:26), or its homologue.

1 28. A pharmaceutical composition for treating vespid venom sensitivity comprising a
 2 peptide of any one of claims 1-27, and a pharmaceutically acceptable carrier.

1 29. A pharmaceutical composition for treating vespid venom sensitivity comprising
 2 more than one peptide of any one of claims 1-27, and a pharmaceutically acceptable carrier.

1 30. A method for treating sensitivity to vespid venom comprising administering to a
2 vespid venom allergic patient a therapeutically effective amount of a peptide of any one of
3 claims 1-27 or the pharmaceutical composition of claim 28 or 29.

1 31. The method according to claim 30 wherein the patient has been identified as
2 sensitive to vespid venom antigen 5.

FIG. 1A

SEQ ID

NO:

1	Ves m	NNYCKIKCLKGGVHTACKYG SLKP	NCGNKKVVSYG LTKQEKQDILKE	1-47
2	Ves v	NNYCKIKCLKGGVHTACKYG SLKP	NCGNKKVVSYG LTKQEKQDILKE	1-47
3	Dol a	NNYCKI CpKG tHTLCKYGTSMKP	NCGgKIVKSYGVTNDEKNEIVKR	1-46
4	Dol mA	NNYCKIKCsrg IHTLCKFGTSMKP	NCGsKIVKvhGVsNDEKNEIVhR	1-47
5	Dol mB	NNYCKIKCrKG IHTLCKFGTSMKP	NCGrnVVKayGITNDEKNEIiKR	1-47
6	Pol a	VDYCKIKCPSG IHTVCQYGESTKPSKNCAGKVIKSVGPTEEEEKLVSE	1-49	
7	Pol e	VDYCKIKCPSG IHTVCQYGESTKPSKNCAGKVIKSVGPTEEEEKLVSE	1-49	

YCKI C G HT C G S KP NC G EK I

1 / 1 1

Ves m	HNDFRQKIARGLETRGNPGPPAKNMKNLVsDELAYiAQVWANQCQY	48-96
Ves v	HNDFRQKIARGLETRGNPGPPAKNMKNLVWDELAYvAQVWANQCQY	48-96
Dol a	HNeFRQKVaqGLETRGNPGPPAKNMNlLVWDELAKIAQTWANQCnF	47-95
Dol mA	HNqFRQKVAKGLETRGNPGPPAKNMNVLWDELAKIAQTWANQCsF	48-96
Dol mB	HNdFRQnVAKGLETRGkPGPPAKNMNVLWDELAKIAQTWANQCdF	48-96
Pol a	HNrFRQKVAQGLETRGNPGPPAASDMNDLVWDELAHIAQVWASQCQFL	50-99
Pol e	HNrFRQKVAQGLETRGNPGPPAASDMNDLVWDELAHIAQVWASQCQFL	50-99

HN FRQ A GLETRG PGQP A M LVW DELA AQ WA QC

FIG. 1B

Ves m	GHDTCRDVAKYQVGQNVALTGSTAAVnDPVKLVKMWEDEVKDYNPKKKF	97-146
Ves v	GHDTCRDVAKYQVGQNVALTGSTAAKYdDPVKLVKMWEDEVKDYNPKKKF	97-146
Dol a	GHDQCRNTAKYpVGQNVAIASSTTGNsYqTMSyLIKMWEDeVKDYNPhKdl	96-145
Dol mA	GHDQCRNTeKYQVGQNVAIASSTTGNsYaTMSKLIeMWEnEVKDFNPKKGt	97-146
Dol mB	nHDdCRNTAKYQVGQNIAlSSTTatqfdrpSKLIKqWEdEVteFNyKvGI	97-146
Pol a	VHDKCRNTAKYPVGQNIa YAGGSnLPDVVSLIKLWENeVKDFNYNTGI	100-147
Pole	VHDKCRNTAKYPVGQNIa YAGGSkLPDVVSLIKLWENeVKDFNYNTGI	100-147

HD	CR	KY	VGQNA	L	WE	EV	N
----	----	----	-------	-------	---	----	----	---	------

2 / 1 1

Ves m	SeNn FLKiGHYtQMvVWANTKEVGCGSiKIYIQEnWHKHylVCNYGpSGNFqNEELYQTK	147-204
Ves v	SgNd FLKtGHYtQMvVWANTKEVGCGSiKIYIQEKWHKHylVCNYGpSGNFmNEELYQTK	147-204
Dol a	mhNN FSKVGHYtQMvWGKTKEIGCGSVKIYIENKWHThylVCNYGpAGNYMNQPvYERK	146-203
Dol mA	igdnN FSKVGHYtQMvWGKTKEIGCGSVKIYIENNWHThylVCNYGpAGNYMdQPIYERK	147-205
Dol mB	qnsN FrKVGHYtQMvWGKTKEIGCGSiKIYIEdNwYThylVCNYGpGndfnQPIYERK	147-204
Pol a	TKQN FAKIGHYtQMvWGKTKEIGCGSLKYmENNMQNHylICNYGPAGNYLGQLPYTKK	148-205
Pole	TKQN FAKIGHYtQMvWGKTKEIGCGSLKYiENkMQNHylICNYGPAGNYLGQLPYTKK	148-205

...	F	K	GHYtQMvW	TKE	GCGS	KY	HYL	CNYGP	GN	Y	K
-----	---	---	----------	-----	------	----	-----	-------	----	---	---

3 / 1 1

FIG.2

SEQ ID NO:

8	1	NNYCKIKCRKGIHTLCKFGT	(1-20)
9	2	GIHTLCKFGTSMKPNCGRNV	(11-30)
10	3	SMKPNCGRNVVKAYGLTNDE	(21-40)
11	4	VKAYGLTNDEKNEILKRHND	(31-50)
12	5	KNEILKRHNDFRQNVAKGLE	(41-60)
13	6	FRQNVAKGLETRGKPGPQPP	(51-70)
14	7	TRGKPGPQPPAKNMNVLVWN	(61-80)
15	8	AKNMNVLVWNDELAKIAQTW	(71-90)
16	9	DELAKIAQTWANQCDFNHDD	(81-100)
17	10	ANQCDFNHDDCRNTAKYQVG	(91-110)
18	11	CRNTAKYQVGQNIAISSTTA	(101-120)
19	12	QNIAISSTTATQFDRPSKLI	(111-130)
20	13	TQFDRPSKLIKQWEDEVTEF	(121-140)
21	14	KQWEDEVTEFNYKVGLQNSN	(131-150)
22	15	NYKVGLQNSNFRKVGHYTQM	(141-160)
23	20	FRKVGHYTQMVWGKT	(151-165)
24	16	HYTQMVGKTKEIGCGSIKY	(156-175)
25	17	KEIGCGSIKYIEDNWYTHYL	(166-185)
26	18	IEDNWYTHYLCNYGPGGND	(176-195)
27	19	VCNYGPGGNDNFNQPIYERK	(186-204)

FIG. 3

<u>SEQ ID</u> <u>NO.</u>					
27	hum	QVQREIVNKHNELRKAV	SPpaSnmLKMEWSr	14-44	4 / 1 1
28	mou	QVQREIVNKHNELRRSV	NPtgSdILKMEWSi	14-44	
29	Ag5	DEKNEILKRHNDFRQNVAKGLETRGKPGPPAKNMNVLVWND		39-81	
	hum	EVTTNQQRWANKCTLQHSDPEDRKTSTRCENL YMSS	DPTSWSSAI	45-89	
	mou	QATTNAQkWANKCILEHSSKDDRKINIRCGENL YMST	DPTLWSTVI	45-89	
	Ag5	ELAKIAQTWANQCDFNHDDCRN TAKYQVGQNIASSTTATQFDRPSKLI		82-130	
	hum	QSWYDEILDVYGVGPKSPN AVVGHYTQLVWYSTYQVGGIAYCPNQDS		90-139	
	mou	QSWYNENEDFVYGVGAK PN SAVGHYTQLVWYSSFKIGCGIAYCPNQDN		90-138	
	Ag5	KQWEDEVTEFNKVGGLQNSNFRKVGHYTQMVVWGKTKEIGCGSIKYIEDNW		131-180	
	hum	LKYYYVCQYCPAGNNMNRKNTPYQQ		140-164	
	mou	LKYFYVCHYCPMGNNVMKKSTPYQQ		139-163	
	Ag5	YTHYLCVNYGPGGNDFNQPIYERK		181-204	

5 / 11

FIG. 4






FIG. 5



6 / 1 1

FIG. 6





7 / 11
FIG. 7

FIG. 8





8 / 11
FIG. 9

FIG. 10





9 / 1 1
FIG. 11

FIG. 12





10/11
FIG. 13

FIG. 14



11 / 11

FIG. 15





FIG. 16



INTERNATIONAL SEARCH REPORT

Intern. al Application No

PCT/US 97/03753

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07K14/435 A61K38/17

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	JOURNAL OF IMMUNOLOGY, vol. 154, no. 2, 15 January 1995, BALTIMORE US, pages 577-584, XP002035178 T.P. KING ET AL.: "Murine T and B Cell Responses to Natural and Recombinant Hornet Venom Allergen Do1 m 5.02 and Its Recombinant Fragments" cited in the application see page 580, right-hand column, paragraph 6 - page 583, right-hand column, paragraph 1; table I --- -/--	1,28,30, 31

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

14 July 1997

Date of mailing of the international search report

24.07.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Fuhr, C

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 97/ 03753

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 30 and 31
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Internat'l Application No
PCT/US 97/03753

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, vol. 97, no. 1, January 1996, page 426 XP002035179 U. MÜLLER ET AL.: "Successful Immunotherapy with T-Cell Epitopes of Bee Venom Phospholipase A2 (PLA) in Patients with Bee Venom (BV) Allergy" see abstract</p> <p style="text-align: center;">---</p>	1,28,30, 31
A	<p>JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, vol. 91, no. 1, January 1993, page 283 XP002035180 T.P. KING ET AL.: "T and B Cell Epitope Mapping with Recombinant Hornet Ag5-2, Dol m 5B" cited in the application see abstract</p> <p style="text-align: center;">---</p>	1,28,30, 31
A	<p>JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, vol. 92, no. 5, November 1993, pages 707-716, XP002035181 D.R. HOFFMAN: "Allergens in Hymenoptera venom XXV: The amino acid sequences of antigen 5 molecules and the structural basis of antigenic cross-reactivity" see page 714, left-hand column, paragraph 2 - page 716, left-hand column, paragraph 1; figures 1-4</p> <p style="text-align: center;">-----</p>	1